arXiv:1512.01510v1 [math.AP] 4 Dec 2015

WELL-POSEDNESS FOR THE CAUCHY PROBLEM OF THE KLEIN-GORDON-ZAKHAROV SYSTEM IN FOUR AND MORE SPATIAL DIMENSIONS

ISAO KATO

Abstract. We study the Cauchy problem for the Klein-Gordon-Zakharov system in spatial dimension $d \geq 4$ with radial or non-radial initial datum $(u, \partial_t u, n, \partial_t n)|_{t=0} \in H^{s+1}(\mathbb{R}^d) \times H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d) \times \dot{H}^{s-1}(\mathbb{R}^d)$. The critical value of s is $s = s_c = d/2 - 2$. If the initial datum is radial, then we prove the small data global well-posedness and scattering at the critical space in $d \geq 4$ by applying the radial Strichartz estimates and U^2, V^2 type spaces. On the other hand, if the initial datum is non-radial, then we prove the local well-posedness at $s = 1/4$ when $d = 4$ and $s = s_c + 1/(d+1)$ when $d \geq 5$ by applying the U^2, V^2 type spaces.

1. Introduction

We consider the Cauchy problem of the Klein-Gordon-Zakharov system:

$$\begin{cases} (\partial_t^2 - \Delta + 1)u = -nu, & (t,x) \in [-T,T] \times \mathbb{R}^d, \\ (\partial_t^2 - c^2\Delta)n = \Delta|u|^2, & (t,x) \in [-T,T] \times \mathbb{R}^d, \\ (u, \partial_t u, n, \partial_t n)|_{t=0} = (u_0, u_1, n_0, n_1) \\ \qquad\qquad \in H^{s+1}(\mathbb{R}^d) \times H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d) \times \dot{H}^{s-1}(\mathbb{R}^d), \end{cases} \tag{1.1}$$

where u, n are real valued functions, $d \geq 4, c > 0$ and $c \neq 1$. The physical model of (1.1) is the interaction of the Langmuir wave and the ion acoustic wave in a plasma. In the physical model, c satisfies $0 < c < 1$. When $d = 3$, Ozawa, Tsutaya and Tsutsumi [26] proved that (1.1) is globally well-posed in the energy space $H^1(\mathbb{R}^3) \times L^2(\mathbb{R}^3) \times L^2(\mathbb{R}^3) \times \dot{H}^{-1}(\mathbb{R}^3)$. They applied the Fourier restriction norm method to obtain the local well-posedness. Then by the local well-posedness and the energy method, they obtained the global well-posedness. For $d = 3$, Guo, Nakanishi and Wang [6] proved scattering in the energy class with small, radial initial data. They applied the normal form reduction and the radial Strichartz estimates. If

2010 *Mathematics Subject Classification.* 35Q55, 35B40, 35A01, 35A02.

Key words and phrases. scattering, well-posedness, Cauchy problem, low regularity, bilinear estimate, radial Strichartz estimate, U^2, V^2 type Bourgain spaces.

we transform $u_\pm := \omega_1 u \pm i\partial_t u, n_\pm := n \pm i(c\omega)^{-1}\partial_t n, \omega_1 := (1-\Delta)^{1/2}, \omega := (-\Delta)^{1/2}$, then (1.1) is equivalent to the following.

$$\begin{cases} (i\partial_t \mp \omega_1)u_\pm = \pm(1/4)(n_+ + n_-)(\omega_1^{-1}u_+ + \omega_1^{-1}u_-), \quad (t,x) \in [-T,T]\times\mathbb{R}^d, \\ (i\partial_t \mp c\omega)n_\pm = \pm(4c)^{-1}\omega|\omega_1^{-1}u_+ + \omega_1^{-1}u_-|^2, \qquad (t,x) \in [-T,T]\times\mathbb{R}^d, \\ (u_\pm, n_\pm)|_{t=0} = (u_{\pm 0}, n_{\pm 0}) \in H^s(\mathbb{R}^d)\times\dot{H}^s(\mathbb{R}^d). \end{cases} \tag{1.2}$$

Our main result is as follows.

Theorem 1.1. *(i) Let $d = 4$. Then (1.2) is locally well-posed in $H^{1/4}(\mathbb{R}^4) \times \dot{H}^{1/4}(\mathbb{R}^4)$.*
(ii) Let $d \geq 5$ and $s = (d^2-3d-2)/2(d+1)$. Then (1.2) is locally well-posed in $H^s(\mathbb{R}^d)\times\dot{H}^s(\mathbb{R}^d)$.
(iii) Let $d \geq 4, s = s_c = d/2-2$ and assume the initial data $(u_{\pm 0}, n_{\pm 0})$ is radial. Then, (1.2) is globally well-posed in $H^s(\mathbb{R}^d)\times\dot{H}^s(\mathbb{R}^d)$.

Corollary 1.2. *The solution obtained in Theorem 1.1 (iii) scatters as $t \to \pm\infty$.*

For more precise statement of Theorem 1.1 and Corollary 1.2, see Propositions 4.1, 4.2. The scaling regularity of (1.2) is $s = s_c = d/2-2$. We consider both the radial case and the non-radial case. First, we consider the radial case. In the radial case, the Strichartz estimates hold for a more wider range of (q,r). More precisely, see Propostions 2.11, 2.12. On the other hand, we have to recover a half derivative loss to derive the key bilinear estimates at the critial space. Thanks to $c > 0$ and $c \neq 1$, if $|\xi| \gg |\xi'|$, then it holds that

$$M' := \max\big\{\big|\tau \pm c|\xi|\big|, |\tau' \pm \langle\xi'\rangle|, |\tau-\tau' \pm \langle\xi-\xi'\rangle|\big\} \gtrsim |\xi|. \tag{1.3}$$

Here, ξ, ξ' denote frequency for the wave equation, Klein-Gordon equation respectively and $\tau \pm c|\xi|$ (resp. $\tau' \pm \langle\xi'\rangle, \tau-\tau' \pm \langle\xi-\xi'\rangle$) denote the symbol of the linear part for the wave equation (resp. the Klein-Gordon equation). From (1.3) and by applying the U^2, V^2 type spaces, then we can recover the derivative loss. Therefore, we can obtain the bilinear estimates at the critical space by applying the radial Strichartz estimates and U^2, V^2 type spaces. Next, we consider $d = 4$ and the non-radial case. When $d \leq 4$, the Lorentz regularity s_l is an important index as well as the scaling regularity for the well-posedness for the wave equation. When $d = 4$ with quadratic nonlinearity, the Lorentz regularity $s_l = 1/4$. On the other hand, $s_c = 0$, so we need to consider $s \geq s_l = 1/4$. When $d = 4$, we obtain local well-posedness at $s = s_l = 1/4$ by applying U^2, V^2 type spaces. Finally, we consider $d \geq 5$ and the

non-radial case. Since $s_c \geq s_l$ when $d \geq 5$, we expect the local well-posedness with $s = s_c$. However, we only obtain the local well-posedness with $s = s_c + 1/(d+1)$. It seems difficult to prove the bilinear estimate with $s = s_c$. The reason is as below. We observe the first equation of (1.2). We regard the nonlinearity as $n_\pm(\omega_1^{-1}u_\pm)$. Here, we consider the following cases. The case $|\xi| \lesssim |\xi'|$ and the case $|\xi| \gg |\xi'|$, where ξ, ξ' denote the frequency of $n_\pm, u_\pm$ respectively. For the case $|\xi| \lesssim |\xi'|$, the nonlinearity does not have the derivative loss, so we can derive the bilinear estimate at the critical space only by applying the Strichartz estimates. However, for the case $|\xi| \gg |\xi'|$, we need to recover a half derivative loss by (1.3). Here, there are three cases in (1.3). The cases $(a)\, M' = \big|\tau \pm c|\xi|\big|$, $(b)\, M' = |\tau' \pm \langle \xi' \rangle|$ and $(c)\, M' = |\tau - \tau' \pm \langle \xi - \xi' \rangle|$. For the case (a) or (c), we apply (1.3) for $n_\pm$ and the Strichartz estimates for $\omega_1^{-1}u_\pm$. Then we can obtain the bilinear estimate at the critical space. Whereas for (b), we apply the Strichartz estimates for $n_\pm$ and apply (1.3) for $\omega_1^{-1}u_\pm$. In this case, we cannot prove the bilinear estimate at the critical space. As a result, we have to impose more regularity.

In section 2, we prepare some notations and lemmas with respect to U^p, V^p, in section 3, we prove the bilinear estimates and in section 4, we prove the main result.

Acknowledgement

The author appreciate Professor K. Tsugawa for giving useful advice. Also, the author would like to thank S. Kinoshita for telling the author about the radial Strichartz estimates for the wave equation.

2. Notations and Preliminary Lemmas

In this section, we prepare some lemmas, propositions and notations to prove the main theorem. $A \lesssim B$ means that there exists $C > 0$ such that $A \leq CB$. Also, $A \sim B$ means $A \lesssim B$ and $B \lesssim A$. Let $u = u(t,x)$. $\mathcal{F}_t u$, $\mathcal{F}_x u$ denote the Fourier transform of u in time, space, respectively. $\mathcal{F}_{t,x} u = \widehat{u}$ denotes the Fourier transform of u in space and time. Let $\mathcal{Z}$ be the set of finite partitions $-\infty < t_0 < t_1 < \cdots < t_K = \infty$ and let $\mathcal{Z}_0$ be the set of finite partitions $-\infty < t_0 < t_1 < \cdots < t_K \leq \infty$.

Definition 1. Let $1 \leq p < \infty$. For $\{t_k\}_{k=0}^K \in \mathcal{Z}$ and $\{\phi_k\}_{k=0}^{K-1} \subset L_x^2$ with $\sum_{k=0}^{K-1} \|\phi_k\|_{L_x^2}^p = 1$, we call the function $a : \mathbb{R} \to L_x^2$ given by

$$a = \sum_{k=1}^{K} \mathbf{1}_{[t_{k-1}, t_k)} \phi_{k-1}$$

a U^p-atom. Furthermore, we define the atomic space

$$U^p := \Big\{ u = \sum_{j=1}^{\infty} \lambda_j a_j \,\Big|\, a_j : U^p\text{-atom}, \lambda_j \in \mathbb{C} \text{ such that } \sum_{j=1}^{\infty} |\lambda_j| < \infty \Big\}$$

with norm

$$\|u\|_{U^p} := \inf\Big\{ \sum_{j=1}^{\infty} |\lambda_j| \,\Big|\, u = \sum_{j=1}^{\infty} \lambda_j a_j, \lambda_j \in \mathbb{C}, a_j : U^p\text{-atom} \Big\}.$$

Proposition 2.1. *Let $1 \le p < q < \infty$.*
(i) U^p is a Banach space.
(ii) The embeddings $U^p \subset U^q \subset L_t^\infty(\mathbb{R}; L_x^2)$ are continuous.
(iii) For $u \in U^p$, it holds that $\lim_{t \to t_0+} \|u(t) - u(t_0)\|_{L_x^2} = 0$, i.e. every $u \in U^p$ is right-continuous.
(iv) The closed subspace U_c^p of all continuous functions in U^p is a Banach space.

The above proposition is in [8] (Proposition 2.2).

Definition 2. Let $1 \le p < \infty$. We define V^p as the normed space of all functions $v : \mathbb{R} \to L_x^2$ such that $\lim_{t \to \pm\infty} v(t)$ exist and for which the norm

$$\|v\|_{V^p} := \sup_{\{t_k\}_{k=0}^K \in \mathcal{Z}} \Big(\sum_{k=1}^{K} \|v(t_k) - v(t_{k-1})\|_{L_x^2}^p \Big)^{1/p}$$

is finite, where we use the convention that $v(-\infty) := \lim_{t \to -\infty} v(t)$ and $v(\infty) := 0$. Likewise, let V_-^p denote the closed subspace of all $v \in V^p$ with $\lim_{t \to -\infty} v(t) = 0$.

The definitions of V^p and V_-^p, see the erratum [9].

Proposition 2.2. *Let $1 \le p < q < \infty$.*
(i) Let $v : \mathbb{R} \to L_x^2$ be such that

$$\|v\|_{V_0^p} := \sup_{\{t_k\}_{k=0}^K \in \mathcal{Z}_0} \Big(\sum_{k=1}^{K} \|v(t_k) - v(t_{k-1})\|_{L_x^2}^p \Big)^{1/p}$$

is finite. Then, it follows that $v(t_0^+) := \lim_{t \to t_0+} v(t)$ exists for all $t_0 \in [-\infty, \infty)$ and $v(t_0^-) := \lim_{t \to t_0-} v(t)$ exists for all $t_0 \in (-\infty, \infty]$ and moreover,

$$\|v\|_{V^p} = \|v\|_{V_0^p}.$$

(ii) We define the closed subspace V_{rc}^p ($V_{-,rc}^p$) of all right-continuous V^p functions (V_-^p functions). The spaces V^p, V_{rc}^p, V_-^p and $V_{-,rc}^p$ are Banach spaces.
(iii) The embeddings $U^p \subset V_{-,rc}^p \subset U^q$ are continuous.
(iv) The embeddings $V^p \subset V^q$ and $V_-^p \subset V_-^q$ are continuous.

The proof of Proposition 2.2 is in [8] (Proposition 2.4 and Corollary 2.6). Let $\{\mathcal{F}_\xi^{-1}[\varphi_n](x)\}_{n\in\mathbb{Z}} \subset \mathcal{S}(\mathbb{R}^d)$ be the Littlewood-Paley decomposition with respect to x, that is to say

$$\begin{cases} \varphi(\xi) \geq 0, \\ \operatorname{supp}\varphi(\xi) = \{\xi \,|\, 2^{-1} \leq |\xi| \leq 2\}, \end{cases}$$

$$\varphi_n(\xi) := \varphi(2^{-n}\xi), \ \sum_{n=-\infty}^{\infty} \varphi_n(\xi) = 1 \ (\,\xi \neq 0), \ \psi(\xi) := 1 - \sum_{n=0}^{\infty} \varphi_n(\xi).$$

Let $N = 2^n$ $(n \in \mathbb{Z})$ be dyadic number. P_N and $P_{<1}$ denote

$$\begin{aligned} &\mathcal{F}_x[P_N f](\xi) := \varphi(\xi/N)\mathcal{F}_x[f](\xi) = \varphi_n(\xi)\mathcal{F}_x[f](\xi), \\ &\mathcal{F}_x[P_{<1} f](\xi) := \psi(\xi)\mathcal{F}_x[f](\xi). \end{aligned}$$

Let $P_{\geq 1} := \sum_{N\geq 1} P_N$. Similarly, let $\tilde{Q}_N$ be

$$\mathcal{F}_t[\tilde{Q}_N g](\tau) := \phi(\tau/N)\mathcal{F}_t[g](\tau) = \phi_n(\tau)\mathcal{F}_t[g](\tau),$$

where $\{\mathcal{F}_\tau^{-1}[\phi_n](t)\}_{n\in\mathbb{Z}} \subset \mathcal{S}(\mathbb{R})$ be the Littlewood-Paley decomposition with respect to t.

Let $K_\pm(t) = \exp\{\mp it(1-\Delta)^{1/2}\} : L_x^2 \to L_x^2$ be the Klein-Gordon unitary operator such that $\mathcal{F}_x[K_\pm(t)u_0](\xi) = \exp\{\mp it\langle\xi\rangle\}\,\mathcal{F}_x[u_0](\xi)$. Similarly, we define the wave unitary operator $W_{\pm c}(t) = \exp\{\mp ict(-\Delta)^{1/2}\} : L_x^2 \to L_x^2$ such that $\mathcal{F}_x[W_{\pm c}(t)n_0](\xi) = \exp\{\mp ict|\xi|\}\,\mathcal{F}_x[n_0](\xi)$.

Definition 3. We define
$(i)\, U^p_{K_\pm} = K_\pm(\cdot)U^p$ with norm $\|u\|_{U^p_{K_\pm}} = \|K_\pm(-\cdot)u\|_{U^p}$,
$(ii)\, V^p_{K_\pm} = K_\pm(\cdot)V^p$ with norm $\|u\|_{V^p_{K_\pm}} = \|K_\pm(-\cdot)u\|_{V^p}$.
For dyadic number N, M,

$$Q_N := K_\pm(\cdot)\tilde{Q}_N K_\pm(-\cdot), \quad Q_{\geq M} := \sum_{N\geq M} Q_N, \quad Q_{<M} := Id - Q_{\geq M}.$$

Here summation over N means that summation over $n \in \mathbb{Z}$. Similarly, we define $U^p_{W_{\pm c}}, V^p_{W_{\pm c}}$.

Remark 2.1. For L_x^2 unitary operator $A = K_\pm$ or $W_{\pm c}$,

$$U^2_A \subset V^2_{-,\,rc,\,A} \subset L^\infty(\mathbb{R}; L_x^2)$$

Definition 4. For the Klein-Gordon equation, we define $\dot{Y}^s_{K_\pm}, \dot{Z}^s_{K_\pm}, Y^s_{K_\pm}, Z^s_{K_\pm}$ as the closure of all $u \in C(\mathbb{R}; H^s_x(\mathbb{R}^d))$ such that

$$\|u\|_{\dot{Y}^s_{K_\pm}} := \Big(\sum_N N^{2s}\|P_N u\|^2_{V^2_{K_\pm}}\Big)^{1/2}, \quad \|u\|_{\dot{Z}^s_{K_\pm}} := \Big(\sum_N N^{2s}\|P_N u\|^2_{U^2_{K_\pm}}\Big)^{1/2},$$

$$\|u\|_{Y^s_{K_\pm}} := \|P_{<1}u\|_{V^2_{K_\pm}} + \Big(\sum_{N\geq 1} N^{2s}\|P_N u\|^2_{V^2_{K_\pm}}\Big)^{1/2},$$

$$\|u\|_{Z^s_{K_\pm}} := \|P_{<1}u\|_{U^2_{K_\pm}} + \Big(\sum_{N\geq 1} N^{2s}\|P_N u\|^2_{U^2_{K_\pm}}\Big)^{1/2}.$$

Similarly, for the wave equation, we define $\dot{Y}^s_{W_{\pm c}}, \dot{Z}^s_{W_{\pm c}}, Y^s_{W_{\pm c}}, Z^s_{W_{\pm c}}$ by replacing $K_\pm$ with $W_{\pm c}$ in the above norms.

Definition 5. For a Hilbert space H and a Banach space $X \subset C(\mathbb{R}; H)$, we define

$$B_r(H) := \{f \in H \,|\, \|f\|_H \leq r\},$$
$$X([0,T)) := \{u \in C([0,T); H) \,|\, {}^\exists\tilde{u} \in X, \tilde{u}(t) = u(t), t \in [0,T)\}$$

endowed with the norm $\|u\|_{X([0,T))} = \inf\{\|\tilde{u}\|_X |\, \tilde{u}(t) = u(t), t \in [0,T)\}$.

We denote the Duhamel term

$$I_{T,K_\pm}(n,v) := \pm\int_0^t \mathbf{1}_{[0,T]}(t')K_\pm(t-t')n(t')(\omega_1^{-1}v(t'))dt',$$
$$I_{T,W_{\pm c}}(u,v) := \pm\int_0^t \mathbf{1}_{[0,T]}(t')W_{\pm c}(t-t')\omega\big((\omega_1^{-1}u(t'))(\overline{\omega_1^{-1}v(t')})\big)dt'$$

for the Klein-Gordon equation and the wave equation respectively.

Lemma 2.3. *Let $c > 0, c \neq 1$ and $\tau_3 = \tau_1 - \tau_2,\ \xi_3 = \xi_1 - \xi_2$. If $|\xi_1| \gg |\xi_2|$ or $|\xi_1| \ll |\xi_2|$, then it holds that*

$$\max\big\{\big|\tau_1 \pm |\xi_1|\big|, \big|\tau_2 \pm |\xi_2|\big|, \big|\tau_3 \pm c|\xi_3|\big|\big\} \gtrsim \max\{|\xi_1|, |\xi_2|\}. \tag{2.1}$$

Proof. We only prove the case $|\xi_1| \gg |\xi_2|$ since the case $|\xi_1| \ll |\xi_2|$ is proved by the same manner.

$$\text{(l.h.s)} \gtrsim |(\tau_1 \pm |\xi_1|) - (\tau_2 \pm |\xi_2|) - (\tau_3 \pm c|\xi_3|)| \tag{2.2}$$

If $0 < c < 1$, then the right hand side of (2.2) is bounded by

$$|\xi_1| - |\xi_2| - c|\xi_1 - \xi_2| \gtrsim |\xi_1|.$$

If $c > 1$, then the right hand side of (2.2) is bounded by

$$c|\xi_1 - \xi_2| - |\xi_1| - |\xi_2| \gtrsim |\xi_1|.$$

□

The following proposition is in [8] (Theorem 2.8 and Proposition 2.10).

Proposition 2.4. *$u \in V_-^1 \subset U^2$ be absolutely continuous on compact intervals and $v \in V^2$. Then, $\|u\|_{U^2} = \sup_{v\in V^2,\, \|v\|_{V^2}=1} \Big| \int_{-\infty}^{\infty} \langle u'(t), v(t)\rangle_{L_x^2} dt \Big|$.*

Corollary 2.5. *Let $A = K_\pm$ or $W_{\pm c}$ and $u \in V^1_{-,\,A} \subset U^2_A$ be absolutely continuous on compact intervals and $v \in V^2_A$. Then,*

$$\|u\|_{U^2_A} = \sup_{v\in V^2_A,\, \|v\|_{V^2_A}=1} \Big| \int_{-\infty}^{\infty} \langle A(t)(A(-\cdot)u)'(t), v(t)\rangle_{L_x^2} dt \Big|.$$

Lemma 2.6. *Let $M > 0$ and $Q \in \{Q_{<M}, Q_{\geq M}\}$. For $1 \leq p \leq \infty$ and $f \in V^2_{K_\pm}$, it holds that*

$$\|Q(\mathbf{1}_{[0,T]} f)\|_{L_t^p L_x^2} \lesssim T^{1/p} \|f\|_{V^2_{K_\pm}}. \tag{2.3}$$

Proof. By scaling, we only prove (2.3) for $M = 1$. We will show (2.3) for $Q = Q_{<1}$, since $Q_{\geq 1} = Id - Q_{<1}$. Put $g := K_\pm(-\cdot) f$. Then (2.3) is equivalent to

$$\|Q_{<1}(\mathbf{1}_{[0,T]}) K_\pm(\cdot) g)\|_{L_t^p L_x^2} \lesssim T^{1/p} \|g\|_{V^2}. \tag{2.4}$$

By the unitarity of $K_\pm$, we have

$$\begin{aligned} \|Q_{<1}(\mathbf{1}_{[0,T]} K_\pm(\cdot) g)\|_{L_t^p L_x^2} &= \Big\| \sum_{N<1} K_\pm(\cdot) \tilde{Q}_N K_\pm(-\cdot)(\mathbf{1}_{[0,T]} K_\pm(\cdot) g) \Big\|_{L_t^p L_x^2} \\ &= \Big\| \sum_{N<1} \tilde{Q}_N (\mathbf{1}_{[0,T]} g) \Big\|_{L_t^p L_x^2} \\ &= \|\tilde{Q}_{<1}(\mathbf{1}_{[0,T]} g)\|_{L_t^p L_x^2}. \end{aligned} \tag{2.5}$$

For some Schwartz function ϕ, it holds that

$$\tilde{Q}_{<1} h = \phi *_t h.$$

Hence by the Young inequality and the Hölder inequality, we have

$$\begin{aligned} \|\tilde{Q}_{<1}(\mathbf{1}_{[0,T]} g)\|_{L_t^p L_x^2} &\lesssim \|\phi\|_{L_t^1} \|\mathbf{1}_{[0,T]} g\|_{L_t^p L_x^2} \\ &\lesssim \|\mathbf{1}_{[0,T]}\|_{L_t^p} \|g\|_{L_t^\infty L_x^2} \\ &\lesssim T^{1/p} \|g\|_{V^2}. \end{aligned} \tag{2.6}$$

Collecting (2.5)–(2.6), we obtain (2.4). □

Proposition 2.7. *Let $T_0 : L_x^2 \times \ldots \times L_x^2 \to L^1_{loc}(\mathbb{R}^d; \mathbb{C})$ be a n-linear operator. Assume that for some $1 \leq p, q \leq \infty$, it holds that*

$$\|T_0(K_\pm(\cdot)\phi_1, \ldots, K_\pm(\cdot)\phi_n)\|_{L_t^p(\mathbb{R}; L_x^q(\mathbb{R}^d))} \lesssim \prod_{i=1}^n \|\phi_i\|_{L_x^2}.$$

Then, there exists $T : U^p_{K_\pm} \times \ldots \times U^p_{K_\pm} \to L^p_t(\mathbb{R}; L^q_x(\mathbb{R}^d))$ *satisfying*

$$\|T(u_1, \ldots, u_n)\|_{L^p_t(\mathbb{R}; L^q_x(\mathbb{R}^d))} \lesssim \prod_{i=1}^n \|u_i\|_{U^p_{K_\pm}},$$

such that $T(u_1, \ldots, u_n)(t)(x) = T_0(u_1(t), \ldots, u_n(t))(x)$ *a.e.*

See Proposition 2.19 in [8] for the proof of the above proposition.

Proposition 2.8. *Let* $2 \le r < \infty, 2/q = d(1/2 - 1/r), s = 1/q - 1/r + 1/2$. *Then it holds that*

$$\|W_{\pm c}(t)f\|_{L^q_t \dot{W}^{-s,r}_x(\mathbb{R}^{1+d})} \lesssim \|f\|_{L^2_x(\mathbb{R}^d)}.$$

For the proof of Proposition 2.8, see [12], [4].

Proposition 2.9. *Let* $2 \le r < \infty, 2/q = d(1/2 - 1/r), s = 1/q - 1/r + 1/2$. *Then, it holds that*

$$\|K_\pm(t)f\|_{L^q_t W^{-s,r}_x(\mathbb{R}^{1+d})} \lesssim \|f\|_{L^2_x(\mathbb{R}^d)}.$$

For the proof of Proposition 2.9, see [22]. Combining Proposition 2.2, Proposition 2.8, Proposition 2.9 and Proposition 2.7, we have the following proposition.

Proposition 2.10. *Let* $2 \le r < \infty, 2/q = d(1/2 - 1/r), s = 1/q - 1/r + 1/2$. *If* $p > q$, *then it holds that*

$$\|f\|_{L^q_t W^{-s,r}_x(\mathbb{R}^{1+d})} \lesssim \|f\|_{V^p_{K_\pm}}, \qquad \|f\|_{L^q_t \dot{W}^{-s,r}_x(\mathbb{R}^{1+d})} \lesssim \|f\|_{V^p_{W_{\pm c}}}.$$

Proposition 2.11. *Let* $d \ge 3$. *Then, for all radial functions* $f \in L^2_x(\mathbb{R}^d)$, *it holds that*

$$\|W_{\pm c}(t)P_N f\|_{L^q_t L^r_x(\mathbb{R}^{1+d})} \lesssim N^{d(1/2-1/r)-1/q}\|f\|_{L^2_x(\mathbb{R}^d)}, \tag{2.7}$$

if and only if

$$(q, r) = (\infty, 2) \quad or \quad 2 \le q \le \infty, \quad 1/q < (d-1)(1/2 - 1/r). \tag{2.8}$$

See Theorem 1.5 (a) in [7] for the proof of Proposition 2.11.

Proposition 2.12. *If* $2d/(d-1) < q \le \infty, N \ge 1$ *and* $f \in L^2_x(\mathbb{R}^d)$ *is radial function, then it holds that*

$$\|K_\pm(t)P_N f\|_{L^q_{t,x}(\mathbb{R}^{1+d})} \lesssim N^{d/2-(d+1)/q}\|f\|_{L^2_x(\mathbb{R}^d)}. \tag{2.9}$$

If $N < 1$ *and* $f \in L^2_x(\mathbb{R}^d)$ *is radial function, then it holds that*

$$\|K_\pm(t)P_N f\|_{L^q_{t,x}(\mathbb{R}^{1+d})} \lesssim N^{d/2-(d+2)/q}\|f\|_{L^2_x(\mathbb{R}^d)}. \tag{2.10}$$

See (3.13) in [7] for the proof of Proposition 2.12.

Proposition 2.13. *Let (q,r) satisfy $2 \le r \le 2(d+1)/(d-1), r \le q, (1/2)(d-1)(1/2-1/r) \le 1/q < (d-1)(1/2-1/r)$ and $N \ge 1$. Then, for all radial function $f \in L^2_x(\mathbb{R}^d)$, it holds that*

$$\|K_\pm(t)P_N f\|_{L^q_t L^r_x(\mathbb{R}^{1+d})} \lesssim N^{d(1/2-1/r)-1/q}\|f\|_{L^2_x(\mathbb{R}^d)}. \tag{2.11}$$

Proof. When $q = r$, (2.11) follows from (2.9). Interpolating $L^q_{t,x}$ with $L^\infty_t L^2_x$, we obtain (2.11). □

Proposition 2.14. *(i) Let $d \ge 3, (q,r)$ satisfy (2.8) in Proposition 2.11 and $s = d(1/2-1/r)-1/q$. If $p > q$, then it holds that*

$$\|u\|_{L^q_t \dot{W}^{-s,r}_x(\mathbb{R}^{1+d})} \lesssim \|u\|_{V^p_{W_{\pm c}}}.$$

(ii) Let (q,r) satisfy the condition in Proposition 2.13. If $p > q$ and $s_1 = d(1/2-1/r)-1/q$, then it holds that

$$\|P_{\ge 1}u\|_{L^q_t \dot{W}^{-s_1,r}_x(\mathbb{R}^{1+d})} \lesssim \|u\|_{V^p_{K_\pm}}. \tag{2.12}$$

(iii) If $p > q$ and $s_2 = d/2-(d+2)/q$, then it holds that

$$\|P_{<1}u\|_{L^q_t \dot{W}^{-s_2,q}_x(\mathbb{R}^{1+d})} \lesssim \|u\|_{V^p_{K_\pm}}. \tag{2.13}$$

Combining Proposition 2.2, Proposition 2.11 and Proposition 2.7, we have Proposition 2.14 (i). Combining Proposition 2.2, Proposition 2.13 and Proposition 2.7, we have Proposition 2.14 (ii). Combining Proposition 2.2, (2.10) and Proposition 2.7, we have Proposition 2.14 (iii).

Proposition 2.15. *(i) Let $T > 0$ and $u \in Y^s_{K_\pm}([0,T]), u(0) = 0$. Then, there exists $0 \le T' \le T$ such that $\|u\|_{Y^s_{K_\pm}([0,T'])} < \varepsilon$.*
(ii) Let $T > 0$ and $n \in \dot{Y}^s_{W_{\pm c}}([0,T]), n(0) = 0$. Then, there exists $0 \le T' \le T$ such that $\|n\|_{\dot{Y}^s_{W_{\pm c}}([0,T'])} < \varepsilon$.

For the proofs of (i) and (ii), see Proposition 2.24 in [8].

Lemma 2.16. *Let $a \ge 0$. Then for $A = K_\pm$ or $W_{\pm c}$, it holds that*

$$\|\langle \nabla_x \rangle^a f\|_{V^2_A} \lesssim \|f\|_{Y^a_A}.$$

Proof. We only prove for $A = K_\pm$ since we can prove similarly for $A = W_{\pm c}$. By L^2_x orthogonality, we have

$$\begin{aligned}
\|\langle\nabla_x\rangle^a f\|^2_{V^2_{K_\pm}} &\lesssim \sup_{\{t_i\}_{i=0}^I\in\mathcal{Z}} \sum_{i=1}^I (\|P_{<1}(K_\pm(-t_i)f(t_i) - K_\pm(-t_{i-1})f(t_{i-1}))\|^2_{L^2_x} \\
&\quad + \sum_{N\geq 1} N^{2a}\|P_N(K_\pm(-t_i)f(t_i) - K_\pm(-t_{i-1})f(t_{i-1}))\|^2_{L^2_x}) \\
&\lesssim \sup_{\{t_i\}_{i=0}^I\in\mathcal{Z}} \sum_{i=1}^I \|K_\pm(-t_i)P_{<1}f(t_i) - K_\pm(-t_{i-1})P_{<1}f(t_{i-1})\|^2_{L^2_x} \\
&\quad + \sum_{N\geq 1} N^{2a} \sup_{\{t_i\}_{i=0}^I\in\mathcal{Z}} \sum_{i=1}^I \|K_\pm(-t_i)P_N f(t_i) - K_\pm(-t_{i-1})P_N f(t_{i-1})\|^2_{L^2_x} \\
&\lesssim \|f\|^2_{Y^a_{K_\pm}}.
\end{aligned}$$

□

Remark 2.2. Similarly, we see

$$\||\nabla_x|^a f\|_{V^2_A} \lesssim \|f\|_{\dot{Y}^a_A}.$$

Proposition 2.17. *It holds that*

$$\begin{aligned}
&\|Q_M u\|_{L^2_{t,x}(\mathbb{R}^{1+d})} \lesssim M^{-1/2}\|u\|_{V^2_{K_\pm}}, \qquad \|Q_{\geq M} u\|_{L^2_{t,x}(\mathbb{R}^{1+d})} \lesssim M^{-1/2}\|u\|_{V^2_{K_\pm}}, \quad (2.14)\\
&\|Q_{<M} u\|_{V^2_{K_\pm}} \lesssim \|u\|_{V^2_{K_\pm}}, \qquad \|Q_{\geq M} u\|_{V^2_{K_\pm}} \lesssim \|u\|_{V^2_{K_\pm}}, \\
&\|Q_{<M} u\|_{U^2_{K_\pm}} \lesssim \|u\|_{U^2_{K_\pm}}, \qquad \|Q_{\geq M} u\|_{U^2_{K_\pm}} \lesssim \|u\|_{U^2_{K_\pm}}.
\end{aligned}$$

The same estimates hold by replacing the Klein-Gordon operator $K_\pm$ by the wave operator $W_{\pm c}$.

Lemma 2.18. *If f, g are measurable functions, then*

$$\int_{\mathbb{R}^{1+d}} f(t,x)\overline{Q_{\geq M} g(t,x)}dxdt = \int_{\mathbb{R}^{1+d}} (Q_{\geq M} f(t,x))\overline{g(t,x)}dxdt.$$

For the proof of Lemma 2.18, see [13].

Lemma 2.19. *Let $\tilde{u}_{N_1} := \mathbf{1}_{[0,T)}P_{N_1}u, \tilde{v}_{N_2} := \mathbf{1}_{[0,T)}P_{N_2}v, \tilde{n}_{N_3} := \mathbf{1}_{[0,T)}P_{N_3}n, Q_1, Q_2 \in \{Q_{<M}, Q_{\geq M}\}, Q_3 \in \{Q_{<M}, Q_{\geq M}\}, s' := (d^2-3d-2)/2(d+1), s_c := d/2-2$. Then the following estimates hold for sufficiently small $T > 0$ if $\theta > 0$, and hold for all $0 < T < \infty$ if $\theta = 0$ or spherically symmetric (u, v, n):*

(i) If $N_3 \lesssim N_2 \sim N_1$, then

$$|I_1| := \Big| \int_{\mathbb{R}^{1+d}} (\omega_1^{-1}\tilde{u}_{N_1})(\overline{\omega_1^{-1}\tilde{v}_{N_2}})(\overline{\omega \tilde{n}_{N_3}}) dxdt \Big| \\ \lesssim T^\theta N_3^s \|u_{N_1}\|_{V^2_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}},$$

when $(\theta, s) = (1/4, 1/4)$ for $d = 4$ and $(\theta, s) = (0, s_c), (1/(d+1), s')$ for $d \geq 5$. Moreover, if (u, v, n) are spherically symmetric, then for $d \geq 4$,

$$|I_1| \lesssim \langle N_2 \rangle^{(d-8)/3} N_3^{(d+4)/6} \|u_{N_1}\|_{V^2_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}}.$$

(ii) $|I_2| := \big|\int_{\mathbb{R}^{1+d}} \tilde{n}(\omega_1^{-1}\tilde{v})(\overline{P_{<1}\tilde{u}}) dxdt\big| \lesssim T^\theta \|n\|_{\dot{Y}^s_{W_{\pm c}}} \|v\|_{Y^s_{K_\pm}} \|P_{<1}u\|_{V^2_{K_\pm}}$,
when $(\theta, s) = (1, 1/4)$ for $d = 4$ and $(\theta, s) = (0, s_c), (1/(d+1), s')$ for $d \geq 5$. Moreover, if (u, v, n) are spherically symmetric, then for $4 \leq d < 5$,

$$|I_2| \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}} \|v\|_{Y^{s_c}_{K_\pm}} \|P_{<1}u\|_{V^2_{K_\pm}}.$$

(iii) If $N_3 \lesssim N_2 \sim N_1$, then

$$|I_3| := \Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big) (\omega_1^{-1}\tilde{v}_{N_2}) \overline{\tilde{u}_{N_1}} dxdt \Big| \lesssim T^\theta \|n\|_{\dot{Y}^s_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}},$$

when $(\theta, s) = (1/4, 1/4)$ for $d = 4$ and $(\theta, s) = (0, s_c), (1/(d+1), s')$ for $d \geq 5$. Moreover, if (u, v, n) are spherically symmetric, then for $4 \leq d < 5$,

$$|I_3| \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|P_{<1}u\|_{V^2_{K_\pm}}.$$

(iv) If $N_2 \ll N_1 \sim N_3, N_1 \geq 1, M = \varepsilon N_1$ and $\varepsilon > 0$ is sufficiently small, then

$$|I_i| \lesssim T^\theta \|n_{N_3}\|_{V^2_{W_{\pm c}}} \|v\|_{Y^s_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}},$$

when $(\theta, s) = (1/4, 1/4)$ for $d = 4, i = 4, 5, 6$ and $(\theta, s) = (0, s_c)$ for $d \geq 4, i = 4, 6$ and $(\theta, s) = (1/(d+1), s')$ for $d \geq 5, i = 4, 5, 6$. Moreover, if (u, v, n) are spherically symmetric, then for $d \geq 4$, it holds that

$$|I_5| \lesssim \|n_{N_3}\|_{V^2_{W_{\pm c}}} \|v\|_{Y^{sc}_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}},$$

where

$$I_4 := \int_{\mathbb{R}^{1+d}} (Q_{\geq M}\tilde{n}_{N_3}) \Big(\sum_{N_2 \ll N_1} Q_2 \omega_1^{-1} \tilde{v}_{N_2} \Big) (\overline{Q_1 \tilde{u}_{N_1}}) dxdt,$$

$$I_5 := \int_{\mathbb{R}^{1+d}} (Q_3\tilde{n}_{N_3}) \Big(\sum_{N_2 \ll N_1} Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2} \Big) (\overline{Q_1 \tilde{u}_{N_1}}) dxdt,$$

$$I_6 := \int_{\mathbb{R}^{1+d}} (Q_3\tilde{n}_{N_3}) \Big(\sum_{N_2 \ll N_1} Q_2 \omega_1^{-1} \tilde{v}_{N_2} \Big) (\overline{Q_{\geq M} \tilde{u}_{N_1}}) dxdt.$$

Proof. We show (i) first. For $f \in V_A^2,\ A \in \{K_\pm, W_{\pm c}\}$, we see

$$\|\mathbf{1}_{[0,T)} f\|_{V_A^2} \lesssim \|f\|_{V_A^2}. \tag{2.15}$$

First, we show $d = 4$. We apply the Hölder inequality, Proposition 2.10, the Sobolev inequality, (2.15) and $N_3 \lesssim N_1 \sim N_2$, then we have

$$\begin{aligned}
|I_1| &\lesssim \|\mathbf{1}_{[0,T)}\|_{L_t^4} \|\omega_1^{-1}\tilde{u}_{N_1}\|_{L_t^{20/3}L_x^{5/2}} \|\omega_1^{-1}\tilde{v}_{N_2}\|_{L_t^{20/3}L_x^{5/2}} \|\omega\tilde{n}_{N_3}\|_{L_t^{20/9}L_x^5} \\
&\lesssim T^{1/4} \|\langle\nabla_x\rangle^{1/4}\omega_1^{-1}\tilde{u}_{N_1}\|_{V_{K_\pm}^2} \|\langle\nabla_x\rangle^{1/4}\omega_1^{-1}\tilde{v}_{N_2}\|_{V_{K_\pm}^2} \||\nabla_x|^{3/4}\omega\tilde{n}_{N_3}\|_{V_{W_{\pm c}}^2} \\
&\lesssim T^{1/4}\langle N_1\rangle^{1/4-1}\|u_{N_1}\|_{V_{K_\pm}^2} \langle N_2\rangle^{1/4-1}\|v_{N_2}\|_{V_{K_\pm}^2} N_3^{3/4+1}\|n_{N_3}\|_{V_{W_{\pm c}}^2} \\
&\lesssim T^{1/4}N_3^{1/4}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2}\|n_{N_3}\|_{V_{W_{\pm c}}^2}.
\end{aligned}$$

For $d \geq 5$, we apply the Hölder inequality to have

$$|I_1| \lesssim \|\omega_1^{-1}\tilde{u}_{N_1}\|_{L_{t,x}^{2(d+1)/(d-1)}} \|\omega_1^{-1}\tilde{v}_{N_2}\|_{L_{t,x}^{2(d+1)/(d-1)}} \|\omega\tilde{n}_{N_3}\|_{L_{t,x}^{(d+1)/2}}. \tag{2.16}$$

We apply Proposition 2.10, (2.15) and the Sobolev inequality, then we have

$$\begin{aligned}
\|\omega_1^{-1}\tilde{f}_N\|_{L_{t,x}^{2(d+1)/(d-1)}} &\lesssim \langle N\rangle^{1/2-1}\|f_N\|_{V_{K_\pm}^2} = \langle N\rangle^{-1/2}\|f_N\|_{V_{K_\pm}^2}, &(2.17)\\
\|\omega\tilde{n}_{N_3}\|_{L_{t,x}^{(d+1)/2}} &\lesssim \||\nabla_x|^{d(d-5)/2(d-1)}\omega\tilde{n}_{N_3}\|_{L_t^{(d+1)/2}L_x^{2(d^2-1)/(d^2-9)}} \\
&\lesssim \||\nabla_x|^{d/2-2}\omega\tilde{n}_{N_3}\|_{V_{W_{\pm c}}^2} &(2.18)\\
&\lesssim N_3^{s_c+1}\|n_{N_3}\|_{V_{W_{\pm c}}^2} &(2.19)
\end{aligned}$$

Collecting (2.16), (2.17), (2.19) and $N_3 \lesssim N_1 \sim N_2$, we obtain

$$|I_1| \lesssim N_3^{s_c}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2}\|n_{N_3}\|_{V_{W_{\pm c}}^2}.$$

In (2.16), if we apply the Hölder inequality, the Sobolev inequality and Proposition 2.10, then we have

$$\begin{aligned}
\|\omega\tilde{n}_{N_3}\|_{L_{t,x}^{(d+1)/2}} &\lesssim \|\mathbf{1}_{[0,T)}\|_{L_t^{d+1}}\|\omega n_{N_3}\|_{L_t^{d+1}L_x^{(d+1)/2}} \\
&\lesssim T^{1/(d+1)}\||\nabla_x|^{d(d^2-4d-1)/2(d^2-1)}\omega n_{N_3}\|_{L_t^{d+1}L_x^{2(d^2-1)/(d^2-5)}} \\
&\lesssim T^{1/(d+1)}\||\nabla_x|^{(d^2-3d-2)/2(d+1)}\omega n_{N_3}\|_{V_{W_{\pm c}}^2} &(2.20)\\
&\lesssim T^{1/(d+1)}N_3^{s'+1}\|n_{N_3}\|_{V_{W_{\pm c}}^2}. &(2.21)
\end{aligned}$$

Collecting (2.16), (2.17), (2.21) and $N_3 \lesssim N_1 \sim N_2$, we obtain

$$|I_1| \lesssim T^{1/(d+1)}N_3^{s'}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2}\|n_{N_3}\|_{V_{W_{\pm c}}^2}.$$

Next, we prove for $d \geq 4$ and (u, v, n) are spherically symmetric functions. We apply the Hölder inequality to have

$$|I_1| \lesssim \|\omega_1^{-1}\tilde{u}_{N_1}\|_{L^3_{t,x}} \|\omega_1^{-1}\tilde{v}_{N_2}\|_{L^3_{t,x}} \|\omega\tilde{n}_{N_3}\|_{L^3_{t,x}}. \tag{2.22}$$

We apply Proposition 2.14, (2.15) and $N_3 \lesssim N_2 \sim N_1$, then we have

$$\begin{aligned}
\|\omega_1^{-1}\tilde{u}_{N_1}\|_{L^3_{t,x}} &\lesssim \||\nabla_x|^{(d-4)/6} P_{<1}\omega_1^{-1}\tilde{u}_{N_1}\|_{V^2_{K_\pm}} + \Big\||\nabla_x|^{(d-2)/6} \sum_{N\geq 1} P_N(\omega_1^{-1}\tilde{u}_{N_1})\Big\|_{V^2_{K_\pm}} \\
&\lesssim \langle N_1\rangle^{(d-2)/6-1}\Big(\|P_{<1}u_{N_1}\|_{V^2_{K_\pm}} + \Big\|\sum_{N\geq 1} P_N u_{N_1}\Big\|_{V^2_{K_\pm}}\Big) \\
&\lesssim \langle N_2\rangle^{(d-8)/6}\|u_{N_1}\|_{V^2_{K_\pm}}, \qquad (2.23) \\
\|\omega_1^{-1}\tilde{v}_{N_2}\|_{L^3_{t,x}} &\lesssim \langle N_2\rangle^{(d-8)/6}\|v_{N_2}\|_{V^2_{K_\pm}}, \qquad (2.24) \\
\|\omega\tilde{n}_{N_3}\|_{L^3_{t,x}} &\lesssim \||\nabla_x|^{(d-2)/6}\omega\tilde{n}_{N_3}\|_{V^2_{W_{\pm c}}} \lesssim N_3^{(d+4)/6}\|n_{N_3}\|_{V^2_{W_{\pm c}}}. \qquad (2.25)
\end{aligned}$$

From (2.22)–(2.25), we obtain

$$|I_1| \lesssim \langle N_2\rangle^{(d-8)/3} N_3^{(d+4)/6}\|u_{N_1}\|_{V^2_{K_\pm}}\|v_{N_2}\|_{V^2_{K_\pm}}\|n_{N_3}\|_{V^2_{W_{\pm c}}}.$$

Next, we prove (ii). For $d = 4$, we apply the Hölder inequality, the Sobolev inequality, Remark 2.1, (2.15), Remark 2.2, discarding ω_1^{-1} and Lemma 2.16, we obtain

$$\begin{aligned}
|I_2| &\lesssim \|\mathbf{1}_{[0,T)}\|_{L^1_t}\|\tilde{n}\|_{L^\infty_t L^{16/7}_x}\|\omega_1^{-1}\tilde{v}\|_{L^\infty_t L^{16/7}_x}\|P_{<1}\tilde{u}\|_{L^\infty_t L^8_x} \\
&\lesssim T\||\nabla_x|^{1/4}\tilde{n}\|_{L^\infty_t L^2_x}\|\langle\nabla_x\rangle^{1/4}\omega_1^{-1}\tilde{v}\|_{L^\infty_t L^2_x}\||\nabla_x|^{3/2}P_{<1}\tilde{u}\|_{L^\infty_t L^2_x} \\
&\lesssim T\||\nabla_x|^{1/4}\tilde{n}\|_{V^2_{W_{\pm c}}}\|\langle\nabla_x\rangle^{1/4}\omega_1^{-1}v\|_{V^2_{K_\pm}}\|P_{<1}u\|_{V^2_{K_\pm}} \\
&\lesssim T\|n\|_{\dot{Y}^{1/4}_{W_{\pm c}}}\|v\|_{Y^{1/4}_{K_\pm}}\|P_{<1}u\|_{V^2_{K_\pm}}.
\end{aligned}$$

For $d \geq 5$, by the Hölder inequality to have

$$|I_2| \lesssim \|\tilde{n}\|_{L^{(d+1)/2}_{t,x}}\|\omega_1^{-1}\tilde{v}\|_{L^{2(d+1)/(d-1)}_{t,x}}\|P_{<1}\tilde{u}\|_{L^{2(d+1)/(d-1)}_{t,x}}. \tag{2.26}$$

From (2.17), (2.18), Remark 2.2 and Lemma 2.16, we obtain

$$\begin{aligned}
\|\tilde{n}\|_{L^{(d+1)/2}_{t,x}} &\lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}}, \qquad (2.27) \\
\|\omega_1^{-1}\tilde{v}\|_{L^{2(d+1)/(d-1)}_{t,x}} &\lesssim \|\langle\nabla_x\rangle^{-1/2}v\|_{V^2_{K_\pm}} \lesssim \|\langle\nabla_x\rangle^{s_c}v\|_{V^2_{K_\pm}} \lesssim \|v\|_{Y^{s_c}_{K_\pm}}, \qquad (2.28) \\
\|P_{<1}\tilde{u}\|_{L^{2(d+1)/(d-1)}_{t,x}} &\lesssim \|\langle\nabla_x\rangle^{1/2}P_{<1}u\|_{V^2_{K_\pm}} \lesssim \|P_{<1}u\|_{V^2_{K_\pm}}. \qquad (2.29)
\end{aligned}$$

Collecting (2.26)–(2.29), we obtain

$$|I_2| \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}}\|v\|_{Y^{s_c}_{K_\pm}}\|P_{<1}u\|_{V^2_{K_\pm}}.$$

Also for $d \geq 5$, from (2.20), Remark 2.2, (2.26), (2.28) and (2.29) to have

$$|I_2| \lesssim T^{1/(d+1)} \|n\|_{\dot{Y}^{s'}_{W_{\pm c}}} \|v\|_{Y^{s'}_{K_\pm}} \|P_{<1}u\|_{V^2_{K_\pm}}.$$

We prove for $4 \leq d < 5$ and (u, v, n) are spherically symmetric functions. Due to the operator $P_{<1}$,

$$\begin{aligned} |I_2| &\lesssim \Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_3 \lesssim 1} \tilde{n}_{N_3} \Big) \Big(\sum_{N_2 < 1} \omega_1^{-1} \tilde{v}_{N_2} \Big) (\overline{P_{<1}\tilde{u}}) dxdt \Big| \\ &\quad + \sum_{N_2 \geq 1} \Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big) (\omega_1^{-1} \tilde{v}_{N_2}) (\overline{P_{<1}\tilde{u}}) dxdt \Big| \\ &=: I_{2,1} + I_{2,2}. \end{aligned}$$

First, we estimate $I_{2,2}$. We apply the Hölder inequality to have

$$|I_{2,2}| \lesssim \sum_{N_2 \geq 1} \Big\| \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{L^3_{t,x}} \|\omega_1^{-1} \tilde{v}_{N_2}\|_{L^3_{t,x}} \|P_{<1}\tilde{u}\|_{L^3_{t,x}}. \tag{2.30}$$

By Proposition 2.14, (2.15), $d < 5$, $N_3 \lesssim N_2 \sim N_1$ and the Cauchy-Schwarz inequality, then we have

$$\begin{aligned} \Big\| \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{L^3_{t,x}} &\lesssim \Big\| |\nabla_x|^{(d-2)/6} \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{V^2_{W_{\pm c}}} \\ &\lesssim \Big(\sum_{N_3 \lesssim N_2} N_3^{2(5-d)/3} \Big)^{1/2} \Big(\sum_{N_3 \lesssim N_2} N_3^{d-4} \|n_{N_3}\|^2_{V^2_{W_{\pm c}}} \Big)^{1/2} \\ &\lesssim N_2^{(5-d)/3} \|n\|_{\dot{Y}^{sc}_{W_{\pm c}}}. \end{aligned} \tag{2.31}$$

From $d \geq 4$, (2.23) and (2.24), we see

$$\|P_{<1}\tilde{u}\|_{L^3_{t,x}} \lesssim \|P_{<1}u\|_{V^2_{K_\pm}}, \qquad \|\omega_1^{-1}\tilde{v}_{N_2}\|_{L^3_{t,x}} \lesssim \langle N_2 \rangle^{(d-8)/6} \|v_{N_2}\|_{V^2_{K_\pm}}. \tag{2.32}$$

Collecting (2.30)–(2.32), $N_2 \geq 1$ and applying the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} |I_{2,2}| &\lesssim \sum_{N_2 \geq 1} N_2^{(2-d)/6} \|n\|_{\dot{Y}^{sc}_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|P_{<1}u\|_{V^2_{K_\pm}} \\ &\lesssim \|n\|_{\dot{Y}^{sc}_{W_{\pm c}}} \|v\|_{Y^{sc}_{W_{\pm c}}} \|P_{<1}u\|_{V^2_{K_\pm}}. \end{aligned} \tag{2.33}$$

Next, we estimate $I_{2,1}$. By the Hölder inequality to have

$$|I_{2,1}| \lesssim \Big\| \sum_{N_3 \lesssim 1} \tilde{n}_{N_3} \Big\|_{L^3_{t,x}} \Big\| \sum_{N_2 < 1} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L^3_{t,x}} \|P_{<1}\tilde{u}\|_{L^3_{t,x}}. \tag{2.34}$$

From $4 \le d < 5$, (2.31), (2.32) and discarding ω_1^{-1}, we see

$$\Big\| \sum_{N_3 \lesssim 1} \tilde{n}_{N_3} \Big\|_{L^3_{t,x}} \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}}, \qquad \Big\| \sum_{N_2<1} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L^3_{t,x}} \lesssim \|P_{<1} v\|_{V^2_{K_\pm}}. \tag{2.35}$$

Collecting (2.32)–(2.35), we have

$$|I_{2,1}| \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}} \|P_{<1} v\|_{V^2_{K_\pm}} \|P_{<1} u\|_{V^2_{K_\pm}} \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}} \|v\|_{Y^{s_c}_{K_\pm}} \|P_{<1} u\|_{V^2_{K_\pm}}. \tag{2.36}$$

From (2.33) and (2.36), we obtain $|I_2| \lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}} \|v\|_{Y^{s_c}_{K_\pm}} \|P_{<1} u\|_{V^2_{K_\pm}}$. We prove (iii) for $d = 4$ below. We apply the Hölder inequality, the Sobolev inequality, Proposition 2.10 and (2.15), then we have

$$\begin{aligned} |I_3| &\lesssim \|\mathbf{1}_{[0,T)}\|_{L^4_t} \Big\| \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{L^{20/9}_t L^5_x} \|\omega_1^{-1} \tilde{v}_{N_2}\|_{L^{20/3}_t L^{5/2}_x} \|\tilde{u}_{N_1}\|_{L^{20/3}_t L^{5/2}_x} \\ &\lesssim T^{1/4} \Big\| |\nabla_x|^{3/4} \sum_{N_3 \lesssim N_2} n_{N_3} \Big\|_{V^2_{W_{\pm c}}} \langle N_2 \rangle^{1/4-1} \|v_{N_2}\|_{V^2_{K_\pm}} \langle N_1 \rangle^{1/4} \|u_{N_1}\|_{V^2_{K_\pm}}. \end{aligned} \tag{2.37}$$

By the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \Big\| |\nabla_x|^{3/4} \sum_{N_3 \lesssim N_2} n_{N_3} \Big\|_{V^2_{W_{\pm c}}} &\lesssim \sum_{N_3 \lesssim N_2} N_3^{3/4} \|n_{N_3}\|_{V^2_{W_{\pm c}}} \\ &\lesssim \Big(\sum_{N_3 \lesssim N_2} N_3 \Big)^{1/2} \Big(\sum_{N_3 \lesssim N_2} N_3^{1/2} \|n_{N_3}\|^2_{V^2_{W_{\pm c}}} \Big)^{1/2} \\ &\lesssim N_2^{1/2} \|n\|_{\dot{Y}^{1/4}_{W_{\pm c}}}. \end{aligned} \tag{2.38}$$

Collecting (2.37), (2.38) and $N_1 \sim N_2$, we obtain

$$|I_3| \lesssim T^{1/4} \|n\|_{\dot{Y}^{1/4}_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}}.$$

We prove for $d \ge 5$. We apply the Hölder inequality, (2.18), (2.17), Remark 2.2 and $N_1 \sim N_2$, then we have

$$\begin{aligned} |I_3| &\lesssim \Big\| \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{L^{(d+1)/2}_{t,x}} \|\omega_1^{-1} \tilde{v}_{N_2}\|_{L^{2(d+1)/(d-1)}_{t,x}} \|\tilde{u}_{N_1}\|_{L^{2(d+1)/(d-1)}_{t,x}} \qquad (2.39) \\ &\lesssim \Big\| |\nabla_x|^{s_c} \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{V^2_{W_{\pm c}}} \langle N_2 \rangle^{-1/2} \|v_{N_2}\|_{V^2_{K_\pm}} \langle N_1 \rangle^{1/2} \|u_{N_1}\|_{V^2_{K_\pm}} \\ &\lesssim \|n\|_{\dot{Y}^{s_c}_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}}. \end{aligned}$$

From (2.20) and Remark 2.2, we have

$$\Big\| \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{L^{(d+1)/2}_{t,x}} \lesssim T^{1/(d+1)} \|n\|_{\dot{Y}^{s'}_{W_{\pm c}}}. \tag{2.40}$$

Collecting (2.17), (2.39) and (2.40), we obtain

$$|I_3| \lesssim T^{1/(d+1)} \|n\|_{\dot{Y}^{s'}_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}}.$$

When $4 \le d < 5$ and (u, v, n) are spherically symmetric functions, we apply the Hölder inequality to have

$$|I_3| \lesssim \Big\| \sum_{N_3 \lesssim N_2} \tilde{n}_{N_3} \Big\|_{L^3_{t,x}} \|\omega_1^{-1} \tilde{v}_{N_2}\|_{L^3_{t,x}} \|\tilde{u}_{N_1}\|_{L^3_{t,x}}. \tag{2.41}$$

From (2.23), (2.24), we see

$$\|\tilde{u}_{N_1}\|_{L^3_{t,x}} \lesssim \langle N_2 \rangle^{(d-2)/6} \|u_{N_1}\|_{V^2_{K_\pm}}, \qquad \|\omega_1^{-1}\tilde{v}_{N_2}\|_{L^3_{t,x}} \lesssim \langle N_2 \rangle^{(d-8)/6} \|v_{N_2}\|_{V^2_{K_\pm}}. \tag{2.42}$$

Collecting (2.41), (2.31) and (2.42), we obtain

$$|I_3| \lesssim \|n\|_{\dot{Y}^{sc}_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}}.$$

We prove (iv). The estimate for I_6 is the same manner as the estimate for I_4, so we only estimate I_4, I_5. First, we estimate I_4 for $d = 4$. We apply the Hölder inequality, Proposition 2.17, the Sobolev inequality, Lemma 2.6, Proposition 2.10, (2.15), $\langle N_1 \rangle \sim N_1 \ge 1$ and Lemma 2.16 to have

$$\begin{aligned}
|I_4| &\lesssim \|Q_{\ge M}\tilde{n}_{N_3}\|_{L^2_{t,x}} \Big\| \sum_{N_2 \ll N_1} Q_2 \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L^4_t L^{16/3}_x} \|Q_1 \tilde{u}_{N_1}\|_{L^4_t L^{16/5}_x} \\
&\lesssim N_1^{-1/2} \|\tilde{n}_{N_3}\|_{V^2_{W_{\pm c}}} \Big\| \langle \nabla_x \rangle^{5/4} \sum_{N_2 \ll N_1} Q_2 \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L^4_t L^2_x} \|\langle \nabla_x \rangle^{1/12} Q_1 \tilde{u}_{N_1}\|_{L^4_t L^3_x} \\
&\lesssim N_1^{-1/2} \|n_{N_3}\|_{V^2_{W_{\pm c}}} T^{1/4} \Big\| \langle \nabla_x \rangle^{5/4} \sum_{N_2 \ll N_1} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{V^2_{K_\pm}} \langle N_1 \rangle^{1/2} \|u_{N_1}\|_{V^2_{K_\pm}} \\
&\lesssim T^{1/4} \|n_{N_3}\|_{V^2_{W_{\pm c}}} \|v\|_{Y^{1/4}_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}}.
\end{aligned}$$

Next, we prove for $d \ge 4$. We apply the Hölder inequality to have

$$|I_4| \lesssim \|Q_{\ge M}\tilde{n}_{N_3}\|_{L^2_{t,x}} \Big\| \sum_{N_2 \ll N_1} Q_2 \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L^{d+1}_{t,x}} \|Q_1 \tilde{u}_{N_1}\|_{L^{2(d+1)/(d-1)}_{t,x}}. \tag{2.43}$$

By Proposition 2.17, Proposition 2.10, (2.17) and (2.15), we have

$$\|Q_{\ge M}\tilde{n}_{N_3}\|_{L^2_{t,x}} \lesssim N_1^{-1/2} \|n_{N_3}\|_{V^2_{W_{\pm c}}}, \tag{2.44}$$

$$\|Q_1 \tilde{u}_{N_1}\|_{L^{2(d+1)/(d-1)}_{t,x}} \lesssim \langle N_1 \rangle^{1/2} \|u_{N_1}\|_{V^2_{K_\pm}}. \tag{2.45}$$

We apply the Sobolev inequality, Proposition 2.10, (2.15) and Lemma 2.16, we have

$$\begin{aligned}\Big\| \sum_{N_2\ll N_1} Q_2\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_{t,x}^{d+1}} &\lesssim \Big\| \langle\nabla_x\rangle^{d(d-3)/2(d-1)} \sum_{N_2\ll N_1} Q_2\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_t^{d+1}L_x^{2(d^2-1)/(d^2-5)}}\\ &\lesssim \Big\| \langle\nabla_x\rangle^{d(d-3)/2(d-1)+1/(d-1)-1} \sum_{N_2\ll N_1} \tilde{v}_{N_2}\Big\|_{V_{K_\pm}^2}\\ &\lesssim \|v\|_{Y_{K_\pm}^{sc}}. \end{aligned} \tag{2.46}$$

Collecting (2.43)–(2.46) and $N_1 \geq 1$, we obtain

$$|I_4| \lesssim \|n_{N_3}\|_{V_{W_{\pm c}}^2}\|v\|_{Y_{K_\pm}^{sc}}\|u_{N_1}\|_{V_{K_\pm}^2}.$$

For $d \geq 5$, in (2.43), if we apply the Sobolev inequality, Lemma 2.6 and Lemma 2.16 to have

$$\begin{aligned}\Big\| \sum_{N_2\ll N_1} Q_2\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_{t,x}^{d+1}} &\lesssim \Big\| \langle\nabla_x\rangle^{d(d-1)/2(d+1)} \sum_{N_2\ll N_1} Q_2\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_t^{d+1}L_x^2}\\ &\lesssim T^{1/(d+1)}\Big\| \langle\nabla_x\rangle^{(d^2-3d-2)/2(d+1)} \sum_{N_2\ll N_1} \tilde{v}_{N_2}\Big\|_{V_{K_\pm}^2}\\ &\lesssim T^{1/(d+1)}\|v\|_{Y_{K_\pm}^{s'}}. \end{aligned} \tag{2.47}$$

Collecting (2.43)–(2.45), (2.47) and $N_1 \geq 1$, we obtain

$$|I_4| \lesssim T^{1/(d+1)}\|n_{N_3}\|_{V_{W_{\pm c}}^2}\|v\|_{Y_{K_\pm}^{s'}}\|u_{N_1}\|_{V_{K_\pm}^2}.$$

Next, we prove I_5. When $d = 4$, by the Hölder inequality, the Sobolev inequality, Lemma 2.6, Proposition 2.10, (2.15), Proposition 2.17, $N_1 \sim N_3$ and Lemma 2.16, we have

$$\begin{aligned}|I_5| &\lesssim \|Q_3\tilde{n}_{N_3}\|_{L_t^4L_x^{16/5}}\Big\| \sum_{N_2\ll N_1} Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_t^2L_x^{16/3}}\|Q_1\tilde{u}_{N_1}\|_{L_t^4L_x^2}\\ &\lesssim \||\nabla_x|^{1/12}Q_3\tilde{n}_{N_3}\|_{L_t^4L_x^3}\Big\|\langle\nabla_x\rangle^{5/4} \sum_{N_2\ll N_1} Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_{t,x}^2}T^{1/4}\|u_{N_1}\|_{V_{K_\pm}^2}\\ &\lesssim N_3^{1/2}\|n_{N_3}\|_{V_{W_{\pm c}}^2}N_1^{-1/2}\Big\|\langle\nabla_x\rangle^{5/4} \sum_{N_2\ll N_1} \omega_1^{-1}\tilde{v}_{N_2}\Big\|_{V_{K_\pm}^2}T^{1/4}\|u_{N_1}\|_{V_{K_\pm}^2}\\ &\lesssim T^{1/4}\|n_{N_3}\|_{V_{W_{\pm c}}^2}\|v\|_{Y_{K_\pm}^{1/4}}\|u_{N_1}\|_{V_{K_\pm}^2}. \end{aligned}$$

For $d \geq 5$, by the Hölder inequality, we have

$$|I_5| \lesssim \|Q_3\tilde{n}_{N_3}\|_{L_{t,x}^{2(d+1)/(d-1)}}\Big\| \sum_{N_2\ll N_1} Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2}\Big\|_{L_t^2L_x^{d+1}}\|Q_1\tilde{u}_{N_1}\|_{L_t^{d+1}L_x^2}. \tag{2.48}$$

From (2.45), $N_3 \sim N_1$ and Lemma 2.6, we have

$$\|Q_3 \tilde{n}_{N_3}\|_{L_{t,x}^{2(d+1)/(d-1)}} \lesssim \langle N_1 \rangle^{1/2} \|n_{N_3}\|_{V_{W_{\pm c}}^2}, \tag{2.49}$$

$$\|Q_1 \tilde{u}_{N_1}\|_{L_t^{d+1} L_x^2} \lesssim T^{1/(d+1)} \|u_{N_1}\|_{V_{K_\pm}^2}. \tag{2.50}$$

We apply the Sobolev inequality, Proposition 2.17, (2.15) and Lemma 2.16, we have

$$\begin{aligned}
\Big\| \sum_{N_2 \ll N_1} Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L_t^2 L_x^{d+1}} &\lesssim \Big\| \langle \nabla_x \rangle^{d(d-1)/2(d+1)} \sum_{N_2 \ll N_1} Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L_{t,x}^2} \\
&\lesssim N_1^{-1/2} \Big\| \langle \nabla_x \rangle^{(d^2-3d-2)/2(d+1)} \sum_{N_2 \ll N_1} v_{N_2} \Big\|_{V_{K_\pm}^2} \\
&\lesssim N_1^{-1/2} \|v\|_{Y_{K_\pm}^{s'}}.
\end{aligned} \tag{2.51}$$

Collecting (2.48)–(2.51) and $N_1 \geq 1$, we obtain

$$|I_5| \lesssim T^{1/(d+1)} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v\|_{Y_{K_\pm}^{s'}} \|u_{N_1}\|_{V_{K_\pm}^2}.$$

Finally, we prove for (u, v, n) are spherically symmetric functions and $d \geq 4$. By the Hölder inequality, Proposition 2.14, (2.15), the Sobolev inequality, $1 \leq N_1 \sim N_3$, Proposition 2.17 and Lemma 2.16, we have

$$\begin{aligned}
|I_5| &\lesssim \|Q_3 \tilde{n}_{N_3}\|_{L_t^4 L_x^{2d/(d-1)}} \Big\| \sum_{N_2 \ll N_1} Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L_t^2 L_x^d} \|Q_1 \tilde{u}_{N_1}\|_{L_t^4 L_x^{2d/(d-1)}} \\
&\lesssim N_3^{1/4} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \Big\| \langle \nabla_x \rangle^{(d-2)/2} \sum_{N_2 \ll N_1} Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{L_{t,x}^2} \langle N_1 \rangle^{1/4} \|u_{N_1}\|_{V_{K_\pm}^2} \\
&\lesssim N_1^{1/2} \|n_{N_3}\|_{V_{W_{\pm c}}^2} N_1^{-1/2} \Big\| \langle \nabla_x \rangle^{(d-2)/2} \sum_{N_2 \ll N_1} \omega_1^{-1} \tilde{v}_{N_2} \Big\|_{V_{K_\pm}^2} \|u_{N_1}\|_{V_{K_\pm}^2} \\
&\lesssim \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v\|_{Y_{K_\pm}^{sc}} \|u_{N_1}\|_{V_{K_\pm}^2}.
\end{aligned}$$

□

3. Bilinear estimates

Proposition 3.1. *Let $(\theta, s) = (1/4, 1/4)$ for $d = 4$ and $(\theta, s) = (1/(d+1), (d^2 - 3d - 2)/2(d+1))$ for $d \geq 5$. For sufficiently small $T > 0$,*

$$\|I_{T,K_\pm}(n, v)\|_{Z_{K_\pm}^s} \lesssim T^\theta \|n\|_{\dot{Y}_{W_{\pm c}}^s} \|v\|_{Y_{K_\pm}^s}, \tag{3.1}$$

$$\|I_{T,W_{\pm c}}(u, v)\|_{\dot{Z}_{W_{\pm c}}^s} \lesssim T^\theta \|u\|_{Y_{K_\pm}^s} \|v\|_{Y_{K_\pm}^s}. \tag{3.2}$$

Moreover, we assume (u, v, n) are spherically symmetric functions. Then for $d \geq 4$ and for all $0 < T < \infty$, (3.1), (3.2) also holds with $(\theta, s) = (0, d/2 - 2)$.

Proof. We denote $\tilde{u}_{N_1} := \mathbf{1}_{[0,T)}P_{N_1}u, \tilde{v}_{N_2} := \mathbf{1}_{[0,T)}P_{N_2}v, \tilde{n}_{N_3} := \mathbf{1}_{[0,T)}P_{N_3}n$. First, we prove (3.1).

$$\|I_{T,K_\pm}(n,v)\|^2_{Z^s_{K_\pm}} \lesssim \sum_{i=0}^{3} J_i$$

where

$$J_0 := \Big\| \int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t')P_{<1}(\tilde{n}(\omega_1^{-1}\tilde{v}))(t')dt' \Big\|^2_{U^2_{K_\pm}},$$

$$J_1 := \sum_{N_1\geq 1} N_1^{2s} \Big\| \int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t') \sum_{N_2\sim N_1} \sum_{N_3 \lesssim N_2} P_{N_1}(\tilde{n}_{N_3}(\omega_1^{-1}\tilde{v}_{N_2}))(t')dt' \Big\|^2_{U^2_{W_\pm}},$$

$$J_2 := \sum_{N_1\geq 1} N_1^{2s} \Big\| \int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t') \sum_{N_2\ll N_1} \sum_{N_3\sim N_1} P_{N_1}(\tilde{n}_{N_3}(\omega_1^{-1}\tilde{v}_{N_2}))(t')dt' \Big\|^2_{U^2_{W_\pm}},$$

$$J_3 := \sum_{N_1\geq 1} N_1^{2s} \Big\| \int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t') \sum_{N_2\gg N_1} \sum_{N_3\sim N_2} P_{N_1}(\tilde{n}_{N_3}(\omega_1^{-1}\tilde{v}_{N_2}))(t')dt' \Big\|^2_{U^2_{W_\pm}}.$$

By Corollary 2.5, we have

$$J_0^{1/2} \lesssim \sup_{\|u\|_{V^2_{K_\pm}}=1} \Big| \int_{\mathbb{R}^{1+d}} \tilde{n}(\omega_1^{-1}\tilde{v})(\overline{P_{<1}\tilde{u}})dxdt \Big|. \tag{3.3}$$

For $d=4$ and $s=1/4$, from (3.3), Lemma 2.19 (ii) and $\|P_{<1}u\|_{V^2_{K_\pm}} \lesssim \|u\|_{V^2_{K_\pm}}$, we obtain

$$J_0^{1/2} \lesssim T\|n\|_{\dot{Y}^{1/4}_{W_{\pm c}}} \|v\|_{Y^{1/4}_{K_\pm}}.$$

We apply Corollary 2.5 to have

$$J_1 \lesssim \sum_{N_1\geq 1} N_1^{2s} \sup_{\|u\|_{V^2_{K_\pm}}=1} \Big| \sum_{N_2\sim N_1} \sum_{N_3 \lesssim N_2} \int_{\mathbb{R}^{1+d}} \tilde{n}_{N_3}(\omega_1^{-1}\tilde{v}_{N_2})\overline{\tilde{u}_{N_1}}dxdt \Big|^2. \tag{3.4}$$

For $d=4$ and $s=1/4$, by (3.4), $N_1\sim N_2$, Lemma 2.19 (iii) and $\|u_{N_1}\|_{V^2_{K_\pm}} \lesssim \|u\|_{V^2_{K_\pm}}$, we have

$$J_1 \lesssim \sum_{N_2 \gtrsim 1} N_2^{1/2}T^{1/2}\|n\|^2_{\dot{Y}^{1/4}_{W_{\pm c}}} \|v_{N_2}\|^2_{V^2_{K_\pm}} \lesssim T^{1/2}\|n\|^2_{\dot{Y}^{1/4}_{W_{\pm c}}} \|v\|^2_{Y^{1/4}_{K_\pm}}.$$

For the estimate of J_2, we take $M=\varepsilon N_1$ for sufficiently small $\varepsilon>0$. Then, from Lemma 2.3, we have

$$\begin{aligned} &Q_{<M}\big((Q_{<M}\tilde{n}_{N_3})(Q_{<M}\omega_1^{-1}\tilde{v}_{N_2})\big) \\ &= Q_{<M}\Big[\mathcal{F}^{-1}\Big(\int_{\tau_1=\tau_2+\tau_3,\,\xi_1=\xi_2+\xi_3} \widehat{(Q_{<M}\tilde{n}_{N_3})}(\tau_3,\xi_3)\widehat{(Q_{<M}\omega_1^{-1}}\tilde{v}_{N_2})(\tau_2,\xi_2)\Big)\Big] = 0 \end{aligned}$$

when $N_1 \gg N_2$. Therefore,

$$\tilde{n}_{N_3}(\omega_1^{-1}\tilde{v}_{N_2}) = \sum_{i=1}^{3} F_i,$$

where

$$F_1 := Q_1\big((Q_{\geq M}\tilde{n}_{N_3})(Q_2\omega_1^{-1}\tilde{v}_{N_2})\big), \qquad F_2 := Q_1\big((Q_3\tilde{n}_{N_3})(Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2})\big),$$
$$F_3 := Q_{\geq M}\big((Q_3\tilde{n}_{N_3})(Q_2\omega_1^{-1}\tilde{v}_{N_2})\big).$$

For the estimate of F_1, we apply Corollary 2.5 to have

$$\sum_{N_1\geq 1} N_1^{2s}\Big\|\int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t')\sum_{N_2\ll N_1}\sum_{N_3\sim N_1} P_{N_1}F_1(t')dt'\Big\|_{U^2_{W_\pm}}^2$$
$$\lesssim \sum_{N_1\geq 1} N_1^{2s}\sup_{\|u\|_{V^2_{K_\pm}}=1}\Big|\sum_{N_2\ll N_1}\sum_{N_3\sim N_1}\int_{\mathbb{R}^{1+d}}(Q_{\geq M}\tilde{n}_{N_3})(Q_2\omega_1^{-1}\tilde{v}_{N_2})(\overline{Q_1\tilde{u}_{N_1}})dxdt\Big|^2. \quad (3.5)$$

For $d=4, s=1/4$, from Lemma 2.19 (iv), $N_3\sim N_1\geq 1$ and $\|u_{N_1}\|_{V^2_{K_\pm}}\lesssim \|u\|_{V^2_{K_\pm}}$, the right-hand side of (3.5) is bounded by

$$T^{1/2}\sum_{N_3\gtrsim 1} N_3^{1/2}\|n_{N_3}\|^2_{V^2_{W_{\pm c}}}\|v\|^2_{Y^{1/4}_{K_\pm}} \lesssim T^{1/2}\|n\|^2_{\dot{Y}^{1/4}_{W_{\pm c}}}\|v\|^2_{Y^{1/4}_{K_\pm}}. \quad (3.6)$$

For the estimate of F_2, we apply Corollary 2.5 to have

$$\sum_{N_1\geq 1} N_1^{2s}\Big\|\int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t')\sum_{N_2\ll N_1}\sum_{N_3\sim N_1} P_{N_1}F_2(t')dt'\Big\|_{U^2_{W_\pm}}^2$$
$$\lesssim \sum_{N_1\geq 1} N_1^{2s}\sup_{\|u\|_{V^2_{K_\pm}}=1}\Big|\sum_{N_2\ll N_1}\sum_{N_3\sim N_1}\int_{\mathbb{R}^{1+d}}(Q_3\tilde{n}_{N_3})(Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2})(\overline{Q_1\tilde{u}_{N_1}})dxdt\Big|^2. \quad (3.7)$$

For $d=4, s=1/4$, we apply Lemma 2.19 (iv), $N_3\sim N_1\geq 1$ and $\|u_{N_1}\|_{V^2_{K_\pm}}\lesssim \|u\|_{V^2_{K_\pm}}$, then the right-hand side of (3.7) is bounded by

$$T^{1/2}\sum_{N_3\gtrsim 1} N_3^{1/2}\|n_{N_3}\|^2_{V^2_{W_{\pm c}}}\|v\|^2_{Y^{1/4}_{K_\pm}} \lesssim T^{1/2}\|n\|^2_{\dot{Y}^{1/4}_{W_{\pm c}}}\|v\|^2_{Y^{1/4}_{K_\pm}}. \quad (3.8)$$

For the estimate for F_3, we apply Corollary 2.5 to have

$$\sum_{N_1\geq 1} N_1^{2s}\Big\|\int_0^t \mathbf{1}_{[0,T)}(t')K_\pm(t-t')\sum_{N_2\ll N_1}\sum_{N_3\sim N_1} P_{N_1}F_3(t')dt'\Big\|_{U^2_{W_\pm}}^2$$
$$\lesssim \sum_{N_1\geq 1} N_1^{2s}\sup_{\|u\|_{V^2_{K_\pm}}=1}\Big|\sum_{N_2\ll N_1}\sum_{N_3\sim N_1}\int_{\mathbb{R}^{1+d}}(Q_3\tilde{n}_{N_3})(Q_2\omega_1^{-1}\tilde{v}_{N_2})(\overline{Q_{\geq M}\tilde{u}_{N_1}})dxdt\Big|^2. \quad (3.9)$$

For $d=4, s=1/4$, we apply Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, then the right-hand side of (3.9) is bounded by

$$T^{1/2} \sum_{N_3 \gtrsim 1} N_3^{1/2} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v\|_{Y_{K_\pm}^{1/4}}^2 \lesssim T^{1/2} \|n\|_{\dot{Y}_{W_{\pm c}}^{1/4}}^2 \|v\|_{Y_{K_\pm}^{1/4}}^2. \tag{3.10}$$

Collecting (3.6), (3.8) and (3.10), we obtain $J_2 \lesssim T^{1/2} \|n\|_{\dot{Y}_{W_{\pm c}}^{1/4}}^2 \|v\|_{Y_{K_\pm}^{1/4}}^2$. By Corollary 2.5 and the triangle inequality to have

$$\begin{aligned} J_3 &\lesssim \sum_{N_1 \geq 1} N_1^{2s} \sup_{\|u\|_{V_{K_\pm}^2}=1} \Big| \sum_{N_2 \gg N_1} \sum_{N_3 \sim N_2} \int_{\mathbb{R}^{1+d}} \tilde{n}_{N_3} (\omega_1^{-1} \tilde{v}_{N_2}) \overline{\tilde{u}_{N_1}} dxdt \Big|^2 \\ &\lesssim \sum_{N_1 \geq 1} N_1^{2s} \Big(\sum_{N_2 \gg N_1} \sum_{N_3 \sim N_2} \sup_{\|u\|_{V_{K_\pm}^2}=1} \Big| \int_{\mathbb{R}^{1+d}} \tilde{n}_{N_3} (\omega_1^{-1} \tilde{v}_{N_2}) \overline{\tilde{u}_{N_1}} dxdt \Big| \Big)^2. \end{aligned} \tag{3.11}$$

By the same manner as the estimate for Lemma 2.19 (iii), for $d=4, s=1/4$, we have

$$\Big| \int_{\mathbb{R}^{1+4}} \tilde{n}_{N_3} (\omega_1^{-1} \tilde{v}_{N_2}) \overline{\tilde{u}_{N_1}} dxdt \Big| \lesssim T^{1/4} N_3^{1/4} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \|u_{N_1}\|_{V_{K_\pm}^2}. \tag{3.12}$$

From (3.12), the right-hand side of (3.11) is bounded by

$$\sum_{N_1 \geq 1} \Big(\sum_{N_2 \gg N_1} \sum_{N_3 \sim N_2} N_1^{1/4} T^{1/4} N_3^{1/4} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \Big)^2.$$

Hence, $\|\cdot\|_{l^2 l^1} \lesssim \|\cdot\|_{l^1 l^2}$ and the Cauchy-Schwarz inequality to have

$$\begin{aligned} J_3^{1/2} &\lesssim \sum_{N_2 \gtrsim 1} \sum_{N_3 \sim N_2} \Big(\sum_{N_1 \ll N_2} N_1^{1/2} T^{1/2} N_3^{1/2} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v_{N_2}\|_{V_{K_\pm}^2}^2 \Big)^{1/2} \\ &\lesssim T^{1/4} \sum_{N_2 \gtrsim 1} \sum_{N_3 \sim N_2} N_2^{1/4} N_3^{1/4} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \\ &\lesssim T^{1/4} \|n\|_{\dot{Y}_{W_{\pm c}}^{1/4}} \|v\|_{Y_{K_\pm}^{1/4}}. \end{aligned}$$

We prove (3.2). By Corollary 2.5, we only need to estimate K_i $(i=1,2,3)$:

$$K_1 := \sum_{N_3} N_3^{2s} \sup_{\|n\|_{V_{W_{\pm c}}^2}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1} \tilde{u}_{N_1}) (\overline{\omega_1^{-1} \tilde{v}_{N_2}}) (\overline{\omega \tilde{n}_{N_3}}) dxdt \Big|^2,$$

$$K_2 := \sum_{N_3} N_3^{2s} \sup_{\|n\|_{V_{W_{\pm c}}^2}=1} \Big| \sum_{N_2 \ll N_3} \sum_{N_1 \sim N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1} \tilde{u}_{N_1}) (\overline{\omega_1^{-1} \tilde{v}_{N_2}}) (\overline{\omega \tilde{n}_{N_3}}) dxdt \Big|^2,$$

$$K_3 := \sum_{N_3} N_3^{2s} \sup_{\|n\|_{V_{W_{\pm c}}^2}=1} \Big| \sum_{N_2 \gtrsim N_3} \sum_{N_1 \sim N_2} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1} \tilde{u}_{N_1}) (\overline{\omega_1^{-1} \tilde{v}_{N_2}}) (\overline{\omega \tilde{n}_{N_3}}) dxdt \Big|^2.$$

First, we estimate K_1. Put $K_1 = K_{1,1} + K_{1,2}$ where

$$K_{1,1} := \sum_{N_3 \lesssim 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1}\tilde{u}_{N_1})(\overline{\omega_1^{-1}\tilde{v}_{N_2}}) \times (\overline{\omega \tilde{n}_{N_3}}) dxdt \Big|^2, \tag{3.13}$$

$$K_{1,2} := \sum_{N_3 \gg 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1}\tilde{u}_{N_1})(\overline{\omega_1^{-1}\tilde{v}_{N_2}})(\overline{\omega \tilde{n}_{N_3}}) dxdt \Big|^2.$$

For $d = 4, s = 1/4$, by the same manner as the estimate for Lemma 2.19 (i) and $N_1 \ll N_3 \lesssim 1$, we find

$$\begin{aligned}
&\Big| \int_{\mathbb{R}^{1+4}} \Big(\sum_{N_1 \ll N_3} \omega_1^{-1}\tilde{u}_{N_1} \Big) (\overline{\omega_1^{-1}\tilde{v}_{N_2}})(\overline{\omega \tilde{n}_{N_3}}) dxdt \Big| \\
&\lesssim \|\mathbf{1}_{[0,T)}\|_{L^4_t} \Big\| \sum_{N_1 \ll N_3} \omega_1^{-1}\tilde{u}_{N_1} \Big\|_{L^{20/3}_t L^{5/2}_x} \|\omega_1^{-1}\tilde{v}_{N_2}\|_{L^{20/3}_t L^{5/2}_x} \|\omega \tilde{n}_{N_3}\|_{L^{20/9}_t L^5_x} \\
&\lesssim T^{1/4} \Big\| \langle \nabla_x \rangle^{1/4} \sum_{N_1 \ll N_3} \omega_1^{-1}\tilde{u}_{N_1} \Big\|_{V^2_{K_\pm}} \| \langle \nabla_x \rangle^{1/4} \omega_1^{-1}\tilde{v}_{N_2}\|_{V^2_{K_\pm}} \| |\nabla_x|^{3/4} \omega \tilde{n}_{N_3}\|_{V^2_{W_{\pm c}}} \\
&\lesssim T^{1/4} \langle N_1 \rangle^{-1} \Big\| \langle \nabla_x \rangle^{1/4} \sum_{N_1 \ll N_3} \tilde{u}_{N_1} \Big\|_{V^2_{K_\pm}} \langle N_2 \rangle^{1/4-1} \|v_{N_2}\|_{V^2_{K_\pm}} N_3^{3/4+1} \|n_{N_3}\|_{V^2_{W_{\pm c}}} \\
&\lesssim T^{1/4} \|u\|_{Y^{1/4}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}}.
\end{aligned}$$

Hence,

$$K_{1,1} \lesssim \sum_{N_2 \lesssim 1} N_2^{1/2} (T^{1/4} \|u\|_{Y^{1/4}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}})^2 \lesssim T^{1/2} \|u\|^2_{Y^{1/4}_{K_\pm}} \|v\|^2_{Y^{1/4}_{K_\pm}}.$$

We take $M = \varepsilon N_2$ for sufficiently small $\varepsilon > 0$. Then, from Lemma 2.3, we have

$$\begin{aligned}
&Q_{<M}\omega_1^{-1}\mathbf{1}_{[0,T)}\big((Q_{<M}\omega_1^{-1}\tilde{v}_{N_2})(Q_{<M}\omega\, \tilde{n}_{N_3})\big) \\
&= Q_{<M}\omega_1^{-1}\mathbf{1}_{[0,T)}\Big[\mathcal{F}^{-1}\Big(\int_{\tau_1=\tau_2+\tau_3,\, \xi_1=\xi_2+\xi_3} (\widehat{Q_{<M}\omega_1^{-1}\tilde{v}_{N_2}})(\tau_2,\xi_2)(\widehat{Q_{<M}\omega\, \tilde{n}_{N_3}})(\tau_3,\xi_3) \Big)\Big] \\
&= 0.
\end{aligned}$$

Therefore,

$$(\omega_1^{-1}\tilde{v}_{N_2})(\omega\, \tilde{n}_{N_3}) = \sum_{i=1}^{3} G_i,$$

where

$$\begin{aligned}
&G_1 := Q_{\geq M}\big((Q_2\omega_1^{-1}\tilde{v}_{N_2})(Q_3\omega\, \tilde{n}_{N_3})\big), \qquad G_2 := Q_1\big((Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2})(Q_3\omega\, \tilde{n}_{N_3})\big), \\
&G_3 := Q_1\big((Q_2\omega_1^{-1}\tilde{v}_{N_2})(Q_{\geq M}\omega\, \tilde{n}_{N_3})\big).
\end{aligned}$$

Hence, it follows that

$$K_{1,2} \le K_{1,2,1} + K_{1,2,2} + K_{1,2,3}$$

where

$$K_{1,2,1} := \sum_{N_3 \gg 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1} \tilde{u}_{N_1}) \overline{G_1} dx dt \Big|^2, \tag{3.14}$$

$$K_{1,2,2} := \sum_{N_3 \gg 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1} \tilde{u}_{N_1}) \overline{G_2} dx dt \Big|^2, \tag{3.15}$$

$$K_{1,2,3} := \sum_{N_3 \gg 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (\omega_1^{-1} \tilde{u}_{N_1}) \overline{G_3} dx dt \Big|^2. \tag{3.16}$$

By Lemma 2.18,

$$\begin{aligned} K_{1,2,1} \lesssim \sum_{N_3 \gg 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_{\ge M} \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) \\ \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big|^2. \end{aligned} \tag{3.17}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 5$, for $d = 4, s = 1/4$, we find

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+4}} \Big(\sum_{N_1 \ll N_3} Q_{\ge M} \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big| \\ &\lesssim T^{1/4} \|u\|_{Y^{1/4}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}}. \end{aligned} \tag{3.18}$$

$N_3 \gg 1, N_2 \sim N_3$ implies $N_2 \ge 1$. Hence, from (3.17) and (3.18), we have

$$\begin{aligned} K_{1,2,1} &\lesssim \sum_{N_3 \gg 1} N_3^{1/2} \sup_{\|n\|_{V^2_{K_\pm}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+4}} (Q_{\ge M} \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big|^2 \\ &\lesssim \sum_{N_2 \ge 1} N_2^{1/2} (T^{1/4} \|u\|_{Y^{1/4}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}})^2 \lesssim T^{1/2} \|u\|^2_{Y^{1/4}_{K_\pm}} \|v\|^2_{Y^{1/4}_{K_\pm}}. \end{aligned}$$

By Lemma 2.18,

$$\begin{aligned} K_{1,2,2} \lesssim \sum_{N_3 \gg 1} N_3^{2s} \sup_{\|n\|_{V^2_{W_{\pm c}}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_1 \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_{\ge M} \omega_1^{-1} \tilde{v}_{N_2}}) \\ \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big|^2. \end{aligned} \tag{3.19}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 6$, for $d = 4, s = 1/4$, we find

$$\begin{aligned}&\Big|\int_{\mathbb{R}^{1+4}}\Big(\sum_{N_1\ll N_3} Q_1\omega_1^{-1}\tilde{u}_{N_1}\Big)(\overline{Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2}})(\overline{Q_3\omega\tilde{n}_{N_3}})dxdt\Big|\\ &\lesssim T^{1/4}\|u\|_{Y^{1/4}_{K_\pm}}\|v_{N_2}\|_{V^2_{K_\pm}}\|n_{N_3}\|_{V^2_{W_{\pm c}}}.\end{aligned}\tag{3.20}$$

$N_3 \gg 1, N_2 \sim N_3$ implies $N_2 \geq 1$. Hence, from (3.19) and (3.20), we have

$$\begin{aligned}K_{1,2,2} \lesssim& \sum_{N_3\gg 1} N_3^{1/2}\sup_{\|n\|_{V^2_{K_\pm}}=1}\Big|\sum_{N_2\sim N_3}\sum_{N_1\ll N_3}\int_{\mathbb{R}^{1+4}}(Q_1\omega_1^{-1}\tilde{u}_{N_1})(\overline{Q_{\geq M}\omega_1^{-1}\tilde{v}_{N_2}})\\ &\times(\overline{Q_3\omega\tilde{n}_{N_3}})dxdt\Big|^2\\ \lesssim& \sum_{N_2\geq 1} N_2^{1/2}(T^{1/4}\|u\|_{Y^{1/4}_{K_\pm}}\|v_{N_2}\|_{V^2_{K_\pm}})^2 \lesssim T^{1/2}\|u\|^2_{Y^{1/4}_{K_\pm}}\|v\|^2_{Y^{1/4}_{K_\pm}}.\end{aligned}$$

By Lemma 2.18,

$$\begin{aligned}K_{1,2,3} \lesssim& \sum_{N_3\gg 1} N_3^{2s}\sup_{\|n\|_{V^2_{W_{\pm c}}}=1}\Big|\sum_{N_2\sim N_3}\sum_{N_1\ll N_3}\int_{\mathbb{R}^{1+d}}(Q_1\omega_1^{-1}\tilde{u}_{N_1})(\overline{Q_2\omega_1^{-1}\tilde{v}_{N_2}})\\ &\times(\overline{Q_{\geq M}\omega\tilde{n}_{N_3}})dxdt\Big|^2.\end{aligned}\tag{3.21}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 4$, for $d = 4, s = 1/4$, we find

$$\begin{aligned}&\Big|\int_{\mathbb{R}^{1+4}}\Big(\sum_{N_1\ll N_3} Q_1\omega_1^{-1}\tilde{u}_{N_1}\Big)(\overline{Q_2\omega_1^{-1}\tilde{v}_{N_2}})(\overline{Q_{\geq M}\omega\tilde{n}_{N_3}})dxdt\Big|\\ &\lesssim T^{1/4}\|u\|_{Y^{1/4}_{K_\pm}}\|v_{N_2}\|_{V^2_{K_\pm}}\|n_{N_3}\|_{V^2_{W_{\pm c}}}.\end{aligned}\tag{3.22}$$

$N_3 \gg 1, N_2 \sim N_3$ implies $N_2 \geq 1$. Hence, from (3.21) and (3.22), we have

$$\begin{aligned}K_{1,2,3} \lesssim& \sum_{N_3\gg 1} N_3^{1/2}\sup_{\|n\|_{V^2_{K_\pm}}=1}\Big|\sum_{N_2\sim N_3}\sum_{N_1\ll N_3}\int_{\mathbb{R}^{1+4}}(Q_1\omega_1^{-1}\tilde{u}_{N_1})(\overline{Q_2\omega_1^{-1}\tilde{v}_{N_2}})\\ &\times(\overline{Q_{\geq M}\omega\tilde{n}_{N_3}})dxdt\Big|^2\\ \lesssim& \sum_{N_2\geq 1} N_2^{1/2}(T^{1/4}\|u\|_{Y^{1/4}_{K_\pm}}\|v_{N_2}\|_{V^2_{K_\pm}})^2 \lesssim T^{1/2}\|u\|^2_{Y^{1/4}_{K_\pm}}\|v\|^2_{Y^{1/4}_{K_\pm}}.\end{aligned}$$

By symmetry, the estimate for K_2 is obtained by the same manner as the estimate for K_1. Hence, we estimate K_3. By the triangle inequality, Lemma 2.19 (i) and the

Cauchy-Schwarz inequality, we have

$$K_3^{1/2} \lesssim \sum_{N_2} \sum_{N_1 \sim N_2} \Big\{ \sum_{N_3 \lesssim N_2} N_3^{2s} \sup_{\|n\|_{V_{K_\pm}^2}=1} \Big| \int_{\mathbb{R}^{1+d}} (\omega_1^{-1}\tilde{u}_{N_1})(\overline{\omega_1^{-1}\tilde{v}_{N_2}})(\overline{\omega \tilde{n}_{N_3}}) dxdt \Big|^2 \Big\}^{1/2}. \tag{3.23}$$

If $d = 4, s = 1/4$, then we apply Lemma 2.19 (i) and the Cauchy-Schwarz inequality, the right-hand side of (3.23) is bounded by

$$\begin{aligned}
&\sum_{N_2} \sum_{N_1 \sim N_2} \Big\{ \sum_{N_3 \lesssim N_2} N_3^{1/2} (T^{1/4} N_3^{1/4} \|u_{N_1}\|_{V_{K_\pm}^2} \|v_{N_2}\|_{V_{K_\pm}^2})^2 \Big\}^{1/2} \\
&\lesssim T^{1/4} \sum_{N_2} \sum_{N_1 \sim N_2} (N_2 \|u_{N_1}\|_{V_{K_\pm}^2}^2 \|v_{N_2}\|_{V_{K_\pm}^2}^2)^{1/2} \\
&\lesssim T^{1/4} \Big(\sum_N N^{1/2} \|u_N\|_{V_{K_\pm}^2}^2 \Big)^{1/2} \Big(\sum_N N^{1/2} \|v_N\|_{V_{K_\pm}^2}^2 \Big)^{1/2}.
\end{aligned}$$

Since

$$\sum_{N<1} N^{1/2} \|u_N\|_{V_{K_\pm}^2}^2 \lesssim \sum_{N<1} N^{1/2} \|P_{<1} u\|_{V_{K_\pm}^2}^2 \lesssim \|P_{<1} u\|_{V_{K_\pm}^2}^2,$$

we obtain $K_3^{1/2} \lesssim T^{1/4} \|u\|_{Y_{K_\pm}^{1/4}} \|v\|_{Y_{K_\pm}^{1/4}}$.

Next, we prove (3.1) for $d \geq 5$ and $s = s' = (d^2 - 3d - 2)/2(d+1)$ by the same manner as the proof for $d = 4, s = 1/4$. From (3.3) and Lemma 2.19 (ii), we have

$$J_0^{1/2} \lesssim T^{1/(d+1)} \|n\|_{\dot{Y}_{W_{\pm c}}^{s'}} \|v\|_{Y_{K_\pm}^{s'}}.$$

By (3.4), $N_1 \sim N_2$, Lemma 2.19 (iii) and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, we have

$$J_1 \lesssim \sum_{N_2 \gtrsim 1} N_2^{2s'} T^{2/(d+1)} \|n\|_{\dot{Y}_{W_{\pm c}}^{s'}}^2 \|v_{N_2}\|_{V_{K_\pm}^2}^2 \lesssim T^{2/(d+1)} \|n\|_{\dot{Y}_{W_{\pm c}}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2.$$

From Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, the right-hand side of (3.5) is bounded by

$$T^{2/(d+1)} \sum_{N_3 \gtrsim 1} N_3^{2s'} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v\|_{Y_{K_\pm}^{s'}}^2 \lesssim T^{2/(d+1)} \|n\|_{\dot{Y}_{W_{\pm c}}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2. \tag{3.24}$$

From Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, the right-hand side of (3.7) is bounded by

$$T^{2/(d+1)} \sum_{N_3 \gtrsim 1} N_3^{2s'} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v\|_{Y_{K_\pm}^{s'}}^2 \lesssim T^{2/(d+1)} \|n\|_{\dot{Y}_{W_{\pm c}}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2. \tag{3.25}$$

From Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V^2_{K_\pm}} \lesssim \|u\|_{V^2_{K_\pm}}$, the right-hand side of (3.9) is bounded by

$$T^{2/(d+1)} \sum_{N_3 \gtrsim 1} N_3^{2s'} \|n_{N_3}\|^2_{V^2_{W_{\pm c}}} \|v\|^2_{Y^{s'}_{K_\pm}} \lesssim T^{2/(d+1)} \|n\|^2_{\dot{Y}^{s'}_{W_{\pm c}}} \|v\|^2_{Y^{s'}_{K_\pm}}. \tag{3.26}$$

Collecting (3.24)–(3.26), we obtain $J_2 \lesssim T^{2/(d+1)} \|n\|^2_{\dot{Y}^{s'}_{W_{\pm c}}} \|v\|^2_{Y^{s'}_{K_\pm}}$. By the same manner as the estimate for Lemma 2.19 (iii), we obtain

$$\Big| \int_{\mathbb{R}^{1+d}} \tilde{n}_{N_3} (\omega_1^{-1} \tilde{v}_{N_2}) \overline{\tilde{u}_{N_1}} dxdt \Big| \lesssim T^{1/(d+1)} N_3^{s'} \|n_{N_3}\|_{V^2_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \|u_{N_1}\|_{V^2_{K_\pm}}. \tag{3.27}$$

From (3.27), the right-hand side of (3.11) is bounded by

$$\sum_{N_1 \geq 1} \Big(\sum_{N_2 \gg N_1} \sum_{N_3 \sim N_2} N_1^{s'} T^{1/(d+1)} N_3^{s'} \|n_{N_3}\|_{V^2_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \Big)^2.$$

Hence, $\|\cdot\|_{l^2 l^1} \lesssim \|\cdot\|_{l^1 l^2}$ and the Cauchy-Schwarz inequality to have

$$\begin{aligned} J_3^{1/2} &\lesssim \sum_{N_2 \gtrsim 1} \sum_{N_3 \sim N_2} \Big(\sum_{N_1 \ll N_2} N_1^{2s'} T^{2/(d+1)} N_3^{2s'} \|n_{N_3}\|^2_{V^2_{W_{\pm c}}} \|v_{N_2}\|^2_{V^2_{K_\pm}} \Big)^{1/2} \\ &\lesssim T^{1/(d+1)} \sum_{N_2 \gtrsim 1} \sum_{N_3 \sim N_2} N_2^{2s'} N_3^{2s'} \|n_{N_3}\|_{V^2_{W_{\pm c}}} \|v_{N_2}\|_{V^2_{K_\pm}} \\ &\lesssim T^{1/(d+1)} \|n\|_{\dot{Y}^{s'}_{W_{\pm c}}} \|v\|_{Y^{s'}_{K_\pm}}. \end{aligned}$$

We prove (3.2) for $d \geq 5$, $s = s' = (d^2 - 3d - 2)/2(d+1)$ by the same manner as the proof for $d = 4$, $s = 1/4$. By the Hölder inequality to have

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{\omega_1^{-1} \tilde{v}_{N_2}}) (\overline{\omega \tilde{n}_{N_3}}) dxdt \Big| \\ &\lesssim \Big\| \sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \Big\|_{L^{2(d+1)/(d-1)}_{t,x}} \|\omega_1^{-1} \tilde{v}_{N_2}\|_{L^{2(d+1)/(d-1)}_{t,x}} \|\omega \tilde{n}_{N_3}\|_{L^{(d+1)/2}_{t,x}}. \end{aligned} \tag{3.28}$$

By Proposition 2.10, $N_1 \ll N_3 \lesssim 1$ and discarding ω_1^{-1} to have

$$\begin{aligned} \Big\| \sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \Big\|_{L^{2(d+1)/(d-1)}_{t,x}} &\lesssim \langle N_1 \rangle^{1/2} \| \sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \|_{V^2_{K_\pm}} \\ &\lesssim \| \sum_{N_1 \ll N_3} \tilde{u}_{N_1} \|_{V^2_{K_\pm}} \\ &\lesssim \|P_{<1} u\|_{V^2_{K_\pm}}. \end{aligned} \tag{3.29}$$

From (3.13), (3.28), (3.29), (2.17) and (2.21), we obtain

$$\begin{aligned} K_{1,1} &\lesssim \sum_{N_2 \lesssim 1} N_2^{2s'} (T^{1/(d+1)} \|P_{<1} u\|_{V_{K_\pm}^2} \|v_{N_2}\|_{V_{K_\pm}^2})^2 \\ &\lesssim T^{2/(d+2)} \|P_{<1} u\|_{V_{K_\pm}^2}^2 \sum_{N_2 \lesssim 1} N_2^{2s'} \|v_{N_2}\|_{V_{K_\pm}^2}^2 \\ &\lesssim T^{2/(d+2)} \|u\|_{Y_{K_\pm}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2 . \end{aligned}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 5$, we see

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} Q_{\geq M} \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big| \\ &\lesssim T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v_{N_2}\|_{V_{K_\pm}^2} \|n_{N_3}\|_{V_{W_{\pm c}}^2}. \end{aligned} \tag{3.30}$$

From (3.14), (3.17) and (3.30), we have

$$\begin{aligned} K_{1,2,1} &\lesssim \sum_{N_3 \gg 1} N_3^{2s'} \sup_{\|n\|_{V_{K_\pm}^2} = 1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_{\geq M} \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big|^2 \\ &\lesssim \sum_{N_2 \geq 1} N_2^{2s'} (T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v_{N_2}\|_{V_{K_\pm}^2})^2 \lesssim T^{2/(d+1)} \|u\|_{Y_{K_\pm}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2 . \end{aligned}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 6$, we see

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} Q_1 \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big| \\ &\lesssim T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v_{N_2}\|_{V_{K_\pm}^2} \|n_{N_3}\|_{V_{W_{\pm c}}^2}. \end{aligned} \tag{3.31}$$

From (3.15), (3.19) and (3.31), we have

$$\begin{aligned} K_{1,2,2} &\lesssim \sum_{N_3 \gg 1} N_3^{2s'} \sup_{\|n\|_{V_{K_\pm}^2} = 1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_1 \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dx dt \Big|^2 \\ &\lesssim \sum_{N_2 \geq 1} N_2^{2s'} (T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v_{N_2}\|_{V_{K_\pm}^2})^2 \lesssim T^{2/(d+1)} \|u\|_{Y_{K_\pm}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2 . \end{aligned}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 4$, we see

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} Q_1 \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_{\geq M} \omega \tilde{n}_{N_3}}) dx dt \Big| \\ &\lesssim T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v_{N_2}\|_{V_{K_\pm}^2} \|n_{N_3}\|_{V_{W_{\pm c}}^2}. \end{aligned} \tag{3.32}$$

From (3.16), (3.21) and (3.32), we have

$$\begin{aligned} K_{1,2,3} &\lesssim \sum_{N_3 \gg 1} N_3^{2s'} \sup_{\|n\|_{V_{K_\pm}^2}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_1 \omega_1^{-1} \tilde{u}_{N_1})(\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_{\geq M} \omega \tilde{n}_{N_3}}) dx dt \Big|^2 \\ &\lesssim \sum_{N_2 \geq 1} N_2^{2s'} (T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v_{N_2}\|_{V_{K_\pm}^2})^2 \lesssim T^{2/(d+1)} \|u\|_{Y_{K_\pm}^{s'}}^2 \|v\|_{Y_{K_\pm}^{s'}}^2 . \end{aligned}$$

By symmetry, the estimate for K_2 is obtained by the same manner as the estimate for K_1. We apply Lemma 2.19 (i) and the Cauchy-Schwarz inequality, the right-hand side of (3.23) is bounded by

$$\begin{aligned} &\sum_{N_2} \sum_{N_1 \sim N_2} \Big\{ \sum_{N_3 \lesssim N_2} N_3^{2s'} (T^{1/(d+1)} N_3^{s'} \|u_{N_1}\|_{V_{K_\pm}^2} \|v_{N_2}\|_{V_{K_\pm}^2})^2 \Big\}^{1/2} \\ &\lesssim T^{1/(d+1)} \sum_{N_2} \sum_{N_1 \sim N_2} (N_2^{4s'} \|u_{N_1}\|_{V_{K_\pm}^2}^2 \|v_{N_2}\|_{V_{K_\pm}^2}^2)^{1/2} \\ &\lesssim T^{1/(d+1)} \Big(\sum_N N^{2s'} \|u_N\|_{V_{K_\pm}^2}^2 \Big)^{1/2} \Big(\sum_N N^{2s'} \|v_N\|_{V_{K_\pm}^2}^2 \Big)^{1/2} . \end{aligned}$$

Since $s' > 0$, we have

$$\sum_{N<1} N^{2s'} \|u_N\|_{V_{K_\pm}^2}^2 \lesssim \sum_{N<1} N^{2s'} \|P_{<1} u\|_{V_{K_\pm}^2}^2 \lesssim \|P_{<1} u\|_{V_{K_\pm}^2}^2 .$$

Thus, we obtain $K_3^{1/2} \lesssim T^{1/(d+1)} \|u\|_{Y_{K_\pm}^{s'}} \|v\|_{Y_{K_\pm}^{s'}}$.

Finally, we prove (3.1) for $d \geq 4, s = s_c = d/2 - 2$ and spherically symmetric functions (u, v, n) by the same manner as the proof of $d = 4, s = 1/4$. From (3.3) and Lemma 2.19 (ii), we obtain

$$J_0^{1/2} \lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}} \|v\|_{Y_{K_\pm}^{s_c}} .$$

By (3.4), $N_1 \sim N_2$, Lemma 2.19 (iii) and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, we have

$$J_1 \lesssim \sum_{N_2 \gtrsim 1} N_2^{2s_c} \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}}^2 \|v_{N_2}\|_{V_{K_\pm}^2}^2 \lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 .$$

From Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, the right-hand side of (3.5) is bounded by

$$\sum_{N_3 \gtrsim 1} N_3^{2s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 \lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 . \tag{3.33}$$

From Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, the right-hand side of (3.7) is bounded by

$$\sum_{N_3 \gtrsim 1} N_3^{2s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 \lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 . \tag{3.34}$$

From Lemma 2.19 (iv), $N_3 \sim N_1 \geq 1$ and $\|u_{N_1}\|_{V_{K_\pm}^2} \lesssim \|u\|_{V_{K_\pm}^2}$, the right-hand side of (3.9) is bounded by

$$\sum_{N_3 \gtrsim 1} N_3^{2s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 \lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2 . \tag{3.35}$$

Collecting (3.33)–(3.35), we have $J_2 \lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}}^2 \|v\|_{Y_{K_\pm}^{s_c}}^2$. By the same manner as the estimate for Lemma 2.19 (iii), we obtain

$$\Big| \int_{\mathbb{R}^{1+d}} \tilde{n}_{N_3} (\omega_1^{-1} \tilde{v}_{N_2}) \overline{\tilde{u}_{N_1}} dxdt \Big| \lesssim N_3^{s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \|u_{N_1}\|_{V_{K_\pm}^2} . \tag{3.36}$$

From (3.36), the right-hand side of (3.11) is bounded by

$$\sum_{N_1 \geq 1} \Big(\sum_{N_2 \gg N_1} \sum_{N_3 \sim N_2} N_1^{s_c} N_3^{s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \Big)^2 .$$

Hence, $\|\cdot\|_{l^2 l^1} \lesssim \|\cdot\|_{l^1 l^2}$ and the Cauchy-Schwarz inequality to have

$$\begin{aligned} J_3^{1/2} &\lesssim \sum_{N_2 \gtrsim 1} \sum_{N_3 \sim N_2} \Big(\sum_{N_1 \ll N_2} N_1^{2s_c} N_3^{2s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2}^2 \|v_{N_2}\|_{V_{K_\pm}^2}^2 \Big)^{1/2} \\ &\lesssim \sum_{N_2 \gtrsim 1} \sum_{N_3 \sim N_2} N_2^{2s_c} N_3^{2s_c} \|n_{N_3}\|_{V_{W_{\pm c}}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \\ &\lesssim \|n\|_{\dot{Y}_{W_{\pm c}}^{s_c}} \|v\|_{Y_{K_\pm}^{s_c}} . \end{aligned}$$

We prove (3.2) for $d \geq 4, s = s_c = d/2 - 2$ and spherically symmetric functions (u, v, n) by the same manner as the proof of $d = 4, s = 1/4$. By the Hölder inequality to have

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{\omega_1^{-1} \tilde{v}_{N_2}}) (\overline{\omega \tilde{n}_{N_3}}) dxdt \Big| \\ &\lesssim \Big\| \sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \Big\|_{L_{t,x}^3} \|\omega_1^{-1} \tilde{v}_{N_2}\|_{L_{t,x}^3} \|\omega \tilde{n}_{N_3}\|_{L_{t,x}^3} . \end{aligned} \tag{3.37}$$

Discarding ω_1^{-1}, then $N_1 \ll N_3 \lesssim 1$ and the same manner as (2.23), we find

$$\Big\| \sum_{N_1 \ll N_3} \omega_1^{-1} \tilde{u}_{N_1} \Big\|_{L_{t,x}^3} \lesssim \langle N_1 \rangle^{(d-2)/6} \Big\| \sum_{N_1 \ll N_3} \tilde{u}_{N_1} \Big\|_{V_{K_\pm}^2} \lesssim \|P_{<1} u\|_{V_{K_\pm}^2} . \tag{3.38}$$

Collecting (3.13), (3.37), (3.38), (2.24), (2.25) and $N_2 \sim N_3 \lesssim 1$, we obtain

$$\begin{aligned} K_{1,1} &\lesssim \sum_{N_2 \lesssim 1} N_2^{2s_c} (\|P_{<1}u\|_{V^2_{K_\pm}} \langle N_2 \rangle^{(d-8)/6} \|v_{N_2}\|_{V^2_{K_\pm}} N_2^{(d+4)/6})^2 \\ &\lesssim \|P_{<1}u\|^2_{V^2_{K_\pm}} \sum_{N_2 \lesssim 1} N_2^{2s_c} \|v_{N_2}\|^2_{V^2_{K_\pm}} \\ &\lesssim \|u\|^2_{Y^{s_c}_{K_\pm}} \|v\|^2_{Y^{s_c}_{K_\pm}}. \end{aligned}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 5$, we obtain

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} Q_{\geq M} \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_3 \omega \tilde{n}_{N_3}}) dxdt \Big| \\ &\lesssim \|u\|_{Y^{s_c}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}}. \end{aligned} \tag{3.39}$$

From (3.14), (3.17) and (3.39), we have

$$\begin{aligned} K_{1,2,1} &\lesssim \sum_{N_3 \gg 1} N_3^{2s_c} \sup_{\|n\|_{V^2_{K_\pm}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_{\geq M} \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dxdt \Big|^2 \\ &\lesssim \sum_{N_2 \geq 1} N_2^{2s_c} (\|u\|_{Y^{s_c}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}})^2 \lesssim \|u\|^2_{Y^{s_c}_{K_\pm}} \|v\|^2_{Y^{s_c}_{K_\pm}}. \end{aligned}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 6$, we obtain

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} Q_1 \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_3 \omega \tilde{n}_{N_3}}) dxdt \Big| \\ &\lesssim \|u\|_{Y^{s_c}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}}. \end{aligned} \tag{3.40}$$

From (3.15), (3.19) and (3.40), we have

$$\begin{aligned} K_{1,2,2} &\lesssim \sum_{N_3 \gg 1} N_3^{2s_c} \sup_{\|n\|_{V^2_{K_\pm}}=1} \Big| \sum_{N_2 \sim N_3} \sum_{N_1 \ll N_3} \int_{\mathbb{R}^{1+d}} (Q_1 \omega_1^{-1} \tilde{u}_{N_1}) (\overline{Q_{\geq M} \omega_1^{-1} \tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_3 \omega \tilde{n}_{N_3}}) dxdt \Big|^2 \\ &\lesssim \sum_{N_2 \geq 1} N_2^{2s_c} (\|u\|_{Y^{s_c}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}})^2 \lesssim \|u\|^2_{Y^{s_c}_{K_\pm}} \|v\|^2_{Y^{s_c}_{K_\pm}}. \end{aligned}$$

By the same manner as the estimate for Lemma 2.19 $(iv), i = 4$, we obtain

$$\begin{aligned} &\Big| \int_{\mathbb{R}^{1+d}} \Big(\sum_{N_1 \ll N_3} Q_1 \omega_1^{-1} \tilde{u}_{N_1} \Big) (\overline{Q_2 \omega_1^{-1} \tilde{v}_{N_2}}) (\overline{Q_{\geq M} \omega \tilde{n}_{N_3}}) dxdt \Big| \\ &\lesssim \|u\|_{Y^{s_c}_{K_\pm}} \|v_{N_2}\|_{V^2_{K_\pm}} \|n_{N_3}\|_{V^2_{W_{\pm c}}}. \end{aligned} \tag{3.41}$$

From (3.16), (3.21) and (3.41), we have

$$\begin{aligned} K_{1,2,3} &\lesssim \sum_{N_3\gg 1} N_3^{2s_c} \sup_{\|n\|_{V_{K_\pm}^2}=1} \Big| \sum_{N_2\sim N_3} \sum_{N_1\ll N_3} \int_{\mathbb{R}^{1+d}} (Q_1\omega_1^{-1}\tilde{u}_{N_1})(\overline{Q_2\omega_1^{-1}\tilde{v}_{N_2}}) \\ &\qquad \times (\overline{Q_{\geq M}\omega\tilde{n}_{N_3}})dxdt\Big|^2 \\ &\lesssim \sum_{N_2\geq 1} N_2^{2s_c}(\|u\|_{Y_{K_\pm}^{sc}}\|v_{N_2}\|_{V_{K_\pm}^2})^2 \lesssim \|u\|_{Y_{K_\pm}^{sc}}^2\|v\|_{Y_{K_\pm}^{sc}}^2. \end{aligned}$$

By symmetry, the estimate for K_2 is obtained by the same manner as the estimate for K_1. For $d=4$, from (3.23), Lemma 2.19 (i) and the Cauchy-Schwarz inequality to have

$$\begin{aligned} K_3^{1/2} &\lesssim \sum_{N_2}\sum_{N_1\sim N_2}\Big\{\sum_{N_3\lesssim N_2}(\langle N_2\rangle^{-4/3}N_3^{4/3}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2})^2\Big\}^{1/2} \\ &\lesssim \sum_{N_2}\sum_{N_1\sim N_2}\big(\langle N_2\rangle^{-8/3}N_2^{8/3}\|u_{N_1}\|_{V_{K_\pm}^2}^2\|v_{N_2}\|_{V_{K_\pm}^2}^2\big)^{1/2} \\ &\lesssim \Big(\sum_N \langle N\rangle^{-4/3}N^{4/3}\|u_N\|_{V_{K_\pm}^2}^2\Big)^{1/2}\Big(\sum_N \langle N\rangle^{-4/3}N^{4/3}\|v_N\|_{V_{K_\pm}^2}^2\Big)^{1/2}. \end{aligned}$$

By $\langle N\rangle^{-4/3}\leq 1$, we have

$$\sum_{N<1}\langle N\rangle^{-4/3}N^{4/3}\|u_N\|_{V_{K_\pm}^2}^2 \lesssim \sum_{N<1}N^{4/3}\|P_{<1}u\|_{V_{K_\pm}^2}^2 \lesssim \|P_{<1}u\|_{V_{K_\pm}^2}^2.$$

Hence, for $d=4$, we obtain

$$K_3^{1/2} \lesssim \|u\|_{Y_{K_\pm}^0}\|v\|_{Y_{K_\pm}^0}. \tag{3.42}$$

For $d>4$, from (3.23) and Lemma 2.19 (i), we have

$$\begin{aligned} K_3^{1/2} &\lesssim \sum_{N_2}\sum_{N_1\sim N_2}\Big\{\sum_{N_3\lesssim N_2}N_3^{2s_c}(\langle N_2\rangle^{(d-8)/3}N_3^{(d+4)/6}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2})^2\Big\}^{1/2} \\ &\lesssim \sum_{N_2}\sum_{N_1\sim N_2}\Big(\sum_{N_3\lesssim N_2}N_3^{4(d-2)/3}\langle N_2\rangle^{2(d-8)/3}\|u_{N_1}\|_{V_{K_\pm}^2}^2\|v_{N_2}\|_{V_{K_\pm}^2}^2\Big)^{1/2} \\ &\lesssim \sum_{N_2}\sum_{N_1\sim N_2}N_2^{2(d-2)/3}\langle N_2\rangle^{(d-8)/3}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2}. \end{aligned} \tag{3.43}$$

For $d\leq 8$, then $\langle N_2\rangle^{(d-8)/3}\leq N_2^{(d-8)/3}$. Hence by (3.43) and the Cauchy-Schwarz inequality, for $4<d\leq 8$, we have

$$\begin{aligned} K_3^{1/2} &\lesssim \sum_{N_2}\sum_{N_1\sim N_2}N_2^{d-4}\|u_{N_1}\|_{V_{K_\pm}^2}\|v_{N_2}\|_{V_{K_\pm}^2} \\ &\lesssim \|u\|_{Y_{K_\pm}^{sc}}\|v\|_{Y_{K_\pm}^{sc}}. \end{aligned} \tag{3.44}$$

For $d > 8$ and $N_2 < 1$, it holds that $\langle N_2 \rangle \lesssim 1$. Hence, by (3.43) to have

$$\begin{aligned} K_3^{1/2} &\lesssim \sum_{N_2<1} \sum_{N_1 \sim N_2} N_2^{2(d-2)/3} \|u_{N_1}\|_{V_{K_\pm}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \\ &\lesssim \|u\|_{Y_{K_\pm}^{sc}} \|P_{<1} v\|_{V_{K_\pm}^2}. \end{aligned} \tag{3.45}$$

For $d > 8$ and $N_2 \geq 1$, it holds that $\langle N_2 \rangle^{(d-8)/3} \sim N_2^{(d-8)/3}$. Thus by (3.23) and the Cauchy-Schwarz inequality to have

$$\begin{aligned} K_3^{1/2} &\lesssim \sum_{N_2} \sum_{N_1 \sim N_2} N_2^{d-4} \|u_{N_1}\|_{V_{K_\pm}^2} \|v_{N_2}\|_{V_{K_\pm}^2} \\ &\lesssim \|u\|_{Y_{K_\pm}^{sc}} \|v\|_{Y_{K_\pm}^{sc}}. \end{aligned} \tag{3.46}$$

Collecting (3.42), (3.44)–(3.46), we obtain $K_3^{1/2} \lesssim \|u\|_{Y_{K_\pm}^{sc}} \|v\|_{Y_{K_\pm}^{sc}}$ for $d \geq 4$.

□

4. The proof of the main theorem

We define

$$u_\pm := \omega_1 u \pm i \partial_t u, \quad n_\pm := n \pm i (c\omega)^{-1} \partial_t n$$

where $\omega_1 := (1 - \Delta)^{1/2}, \omega := (-\Delta)^{1/2}$. Then the wave equation in (1.1) is rewritten into

$$\begin{cases} i\partial_t u_\pm \mp \omega_1 u_\pm = \pm (1/4)(n_+ + n_-)(\omega_1^{-1} u_+ + \omega_1^{-1} u_-), & (t,x) \in [-T,T] \times \mathbb{R}^d, \\ i\partial_t n_\pm \mp c\omega n_\pm = \pm (4c)^{-1} \omega |\omega_1^{-1} u_+ + \omega_1^{-1} u_-|^2, & (t,x) \in [-T,T] \times \mathbb{R}^d, \\ (u_\pm, n_\pm)|_{t=0} = (u_{\pm 0}, n_{\pm 0}) \in H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d). \end{cases} \tag{4.1}$$

Hence by the Duhamel principle, we consider the following integral equation corresponding to (4.1) on the time interval $[0, T)$ with $0 < T \leq \infty$:

$$u_\pm = \Phi_1(u_\pm, n_+, n_-), \quad n_\pm = \Phi_2(n_\pm, u_+, u_-), \tag{4.2}$$

where

$$\begin{aligned} \Phi_1(u_\pm, n_+, n_-) &:= K_\pm(t) u_{\pm 0} \pm (1/4) \{ I_{T,K_\pm}(n_+, u_+)(t) + I_{T,K_\pm}(n_+, u_-)(t) \\ &\quad + I_{T,K_\pm}(n_-, u_+)(t) + I_{T,K_\pm}(n_-, u_-)(t) \}, \\ \Phi_2(n_\pm, u_+, u_-) &:= W_{\pm c}(t) n_{\pm 0} \pm (4c)^{-1} \{ I_{T,W_{\pm c}}(u_+, u_+)(t) + I_{T,W_{\pm c}}(u_+, u_-)(t) \\ &\quad + I_{T,W_{\pm c}}(u_-, u_+)(t) + I_{T,W_{\pm c}}(u_-, u_-)(t) \}. \end{aligned}$$

Proposition 4.1. *(i) Let $s = 1/4$ for $d = 4$ or $s = (d^2 - 3d - 2)/2(d+1)$ for $d \geq 5$. Let $\delta > 0$ be arbitrary. Then, for any initial data $(u_{\pm 0}, n_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d))$, there exists $T > 0$ and a unique solution of (4.2) on $[0, T]$ such that*

$$(u_\pm, n_\pm) \in Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T]) \subset C([0,T]; H^s(\mathbb{R}^d)) \times C([0,T]; \dot{H}^s(\mathbb{R}^d)).$$

Moreover, let $d \geq 4, s = s_c = d/2 - 2$ and $\delta > 0$ be sufficiently small. If $(u_{\pm 0}, n_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d))$ be radial, then for all $0 < T < \infty$, there exists a unique spherically symmetric solution of (4.2) on $[0, T]$ such that

$$(u_\pm, n_\pm) \in Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T]) \subset C([0,T]; H^s(\mathbb{R}^d)) \times C([0,T]; \dot{H}^s(\mathbb{R}^d)).$$

(ii) The flow map obtained by (i):
$B_\delta(H^s(\mathbb{R}^d)) \times B_\delta(\dot{H}^s(\mathbb{R}^d)) \ni (u_{\pm 0}, n_{\pm 0}) \mapsto (u_\pm, n_\pm) \in Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T])$ is Lipschitz continuous.

Remark 4.1. Due to the time reversibility of the Klein-Gordon-Zakharov equation, Porpositions 4.1 also holds in corresponding time interval $[-T, 0]$

Remark 4.2. By (i) in Proposition 4.1 and Remark 4.1, for any $T > 0$, we have solutions to (4.2) $(u_\pm(t), n_\pm(t))$ on $[0, T]$ and $[-T, 0]$. If radial initial data $(u_{\pm 0}, n_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d))$, then we can take T arbitrary large and by uniqueness, spherically symmetric function $(u_\pm(t), n_\pm(t)) \in C((-\infty, \infty); H^s(\mathbb{R}^d)) \times C((-\infty, \infty); \dot{H}^s(\mathbb{R}^d))$ can be defined uniquely.

Proposition 4.2. *Let the spherically symmetric solution $(u_\pm(t), n_\pm(t))$ to (4.2) on $(-\infty, \infty)$ obtained by Proposirion 4.1, Remark 4.1 and Remark 4.2 with radial initial data $(u_{\pm 0}, n_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d))$. Then, there exist $(u_{\pm,+\infty}, n_{\pm,+\infty})$ and $(u_{\pm,-\infty}, n_{\pm,-\infty})$ in $H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d)$ such that*

$$\|u_\pm(t) - K_\pm(t) u_{\pm,+\infty}\|_{H^s_x(\mathbb{R}^d)} + \|n_\pm(t) - W_{\pm c}(t) n_{\pm,+\infty}\|_{\dot{H}^s_x(\mathbb{R}^d)} \to 0$$

as $t \to +\infty$ and

$$\|u_\pm(t) - K_\pm(t) u_{\pm,-\infty}\|_{H^s_x(\mathbb{R}^d)} + \|n_\pm(t) - W_{\pm c}(t) n_{\pm,-\infty}\|_{\dot{H}^s_x(\mathbb{R}^d)} \to 0$$

as $t \to -\infty$.

proof of Proposition 4.1. First, we prove (i). By Proposition 2.10, there exists $C > 0$ such that

$$\|K_\pm(t) u_{\pm 0}\|_{Y^s_{K_\pm}} \leq C \|u_{\pm 0}\|_{H^s}, \qquad \|W_{\pm c}(t) n_{\pm 0}\|_{\dot{Y}^s_{W_{\pm c}}} \leq C \|n_{\pm 0}\|_{\dot{H}^s}.$$

We denote time interval $I := [0,T]$. If $(u_{\pm 0}, n_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d)\times \dot{H}^s(\mathbb{R}^d))$, $(u_\pm, n_\pm) \in B_r(Y^s_{K_\pm}(I) \times \dot{Y}^s_{W_{\pm c}}(I))$, then by Proposition 3.1, for $(\theta, s) = (1/4, 1/4), d = 4$ or for $(\theta, s) = (1/(d+1), (d^2-3d-2)/2(d+1)), d \geq 5$, it holds that

$$
\begin{aligned}
&\|\Phi_1(u_\pm, n_+, n_-)\|_{Y^s_{K_\pm}(I)} \\
&\leq C\|u_{\pm 0}\|_{H^s} + (1/4)CT^\theta(\|n_+\|_{\dot{Y}^s_{W_{+c}}(I)}\|u_+\|_{Y^s_{K_+}(I)} + \|n_+\|_{\dot{Y}^s_{W_{+c}}(I)}\|u_-\|_{Y^s_{K_-}(I)} \\
&\qquad + \|n_-\|_{\dot{Y}^s_{W_{-c}}(I)}\|u_+\|_{Y^s_{K_+}(I)} + \|n_-\|_{\dot{Y}^s_{W_{-c}}(I)}\|u_-\|_{Y^s_{K_-}(I)}) \\
&\leq C\delta + CT^\theta r^2, \\
&\|\Phi_2(n_\pm, u_+, u_-)\|_{\dot{Y}^s_{W_{\pm c}}(I)} \\
&\leq C\|n_{\pm 0}\|_{H^s} + (CT^\theta/4c)(\|u_+\|^2_{Y^s_{K_+}(I)} + 2\|u_+\|_{Y^s_{K_+}(I)}\|u_-\|_{Y^s_{K_-}(I)} + \|u_-\|^2_{Y^s_{K_-}(I)}) \\
&\leq C\delta + CT^\theta r^2/c.
\end{aligned}
$$

We take $r = 2C\delta$ and $T > 0$ satisfying

$$
4CT^\theta r \leq \min\{1, c\}. \tag{4.3}
$$

Then we have

$$
\|\Phi_1(u_\pm, n_+, n_-)\|_{Y^s_{K_\pm}(I)} \leq r, \qquad \|\Phi_2(n_\pm, u_+, u_-)\|_{\dot{Y}^s_{W_{\pm c}}(I)} \leq r.
$$

Hence, (Φ_1, Φ_2) is a map from $B_r(Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T]))$ into itself. Similarly, we assume $(v_{\pm 0}, m_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d))$, $(v_\pm, m_\pm) \in B_r(Y^s_{K_\pm}(I) \times \dot{Y}^s_{W_{\pm c}}(I))$, then it holds that

$$
\begin{aligned}
&\|\Phi_1(u_\pm, n_+, n_-) - \Phi_1(v_\pm, m_+, m_-)\|_{Y^s_{K_\pm}(I)} \\
&\leq (1/4)(\|I_{T,K_\pm}(n_+, u_+)(t) - I_{T,K_\pm}(m_+, v_+)(t)\|_{Y^s_{K_\pm}(I)} \\
&\qquad + \|I_{T,K_\pm}(n_+, u_-)(t) - I_{T,K_\pm}(m_+, v_-)(t)\|_{Y^s_{K_\pm}(I)} \\
&\qquad + \|I_{T,K_\pm}(n_-, u_+)(t) - I_{T,K_\pm}(m_-, v_+)(t)\|_{Y^s_{K_\pm}(I)} \\
&\qquad + \|I_{T,K_\pm}(n_-, u_-)(t) - I_{T,K_\pm}(m_-, v_-)(t)\|_{Y^s_{K_\pm}(I)}).
\end{aligned} \tag{4.4}
$$

By Proposition 3.1, we have

$$
\begin{aligned}
&\|I_{T,K_\pm}(n_+, u_+)(t) - I_{T,K_\pm}(m_+, v_+)(t)\|_{Y^s_{K_\pm}(I)} \\
&\leq CT^\theta(\|n_+ - m_+\|_{\dot{Y}^s_{W_{+c}}(I)}\|u_+\|_{Y^s_{K_+}(I)} + \|m_+\|_{Y^s_{W_{+c}}(I)}\|u_+ - v_+\|_{Y^s_{K_+}(I)}).
\end{aligned} \tag{4.5}
$$

Similarly, we have

$$\begin{aligned}
&\|I_{T,K_\pm}(n_+,u_-)(t)-I_{T,K_\pm}(m_+,v_-)(t)\|_{Y^s_{K_\pm}(I)}\\
&\le CT^\theta(\|n_+-m_+\|_{\dot Y^s_{W_{+c}}(I)}\|u_-\|_{Y^s_{K_-}(I)}+\|m_+\|_{\dot Y^s_{W_{+c}}(I)}\|u_--v_-\|_{Y^s_{K_-}(I)}), \qquad (4.6)\\
&\|I_{T,K_\pm}(n_-,u_+)(t)-I_{T,K_\pm}(m_-,v_+)(t)\|_{Y^s_{K_\pm}(I)}\\
&\le CT^\theta(\|n_--m_-\|_{\dot Y^s_{W_{-c}}(I)}\|u_+\|_{Y^s_{K_+}(I)}+\|m_-\|_{\dot Y^s_{W_{-c}}(I)}\|u_+-v_+\|_{Y^s_{K_+}(I)}), \qquad (4.7)\\
&\|I_{T,K_\pm}(n_-,u_-)(t)-I_{T,K_\pm}(m_-,v_-)(t)\|_{Y^s_{K_\pm}(I)}\\
&\le CT^\theta(\|n_--m_-\|_{\dot Y^s_{W_{-c}}(I)}\|u_-\|_{Y^s_{K_-}(I)}+\|m_-\|_{\dot Y^s_{W_{-c}}(I)}\|u_--v_-\|_{Y^s_{K_-}(I)}). \qquad (4.8)
\end{aligned}$$

Hence from $\|u_\pm\|_{Y^s_{K_\pm}(I)}\le r, \|m_\pm\|_{\dot Y^s_{W_{\pm c}}(I)}\le r$, (4.4)–(4.8) and (4.3), we have

$$\begin{aligned}
&\|\Phi_1(u_\pm,n_+,n_-)-\Phi_1(v_\pm,m_+,m_-)\|_{Y^s_{K_\pm}(I)}\\
&\le (1/8)(\|u_+-v_+\|_{Y^s_{K_+}(I)}+\|u_--v_-\|_{Y^s_{K_-}(I)}\\
&\qquad +\|n_+-m_+\|_{\dot Y^s_{W_{+c}}(I)}+\|n_--m_-\|_{\dot Y^s_{W_{-c}}(I)}). \qquad (4.9)
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
&\|\Phi_2(n_\pm,u_+,u-)-\Phi_2(m_\pm,v_+,v_-)\|_{\dot Y^s_{W_{\pm c}}(I)}\\
&=(4c)^{-1}(\|I_{T,W_{\pm c}}(u_+,u_+)(t)-I_{T,W_{\pm c}}(v_+,v_+)(t)\|_{\dot Y^s_{W_{\pm c}}(I)}\\
&\qquad +\|I_{T,W_{\pm c}}(u_+,u_-)(t)-I_{T,W_{\pm c}}(v_+,v_-)(t)\|_{\dot Y^s_{W_{\pm c}}(I)}\\
&\qquad +\|I_{T,W_{\pm c}}(u_-,u_+)(t)-I_{T,W_{\pm c}}(v_-,v_+)(t)\|_{\dot Y^s_{W_{\pm c}}(I)}\\
&\qquad +\|I_{T,W_{\pm c}}(u_-,u_-)(t)-I_{T,W_{\pm c}}(v_-,v_-)(t)\|_{\dot Y^s_{W_{\pm c}}(I)}). \qquad (4.10)
\end{aligned}$$

By Proposition 3.1, we have

$$\begin{aligned}
&\|I_{T,W_{\pm c}}(u_+,u_+)(t)-I_{T,W_{\pm c}}(v_+,v_+)(t)\|_{\dot Y^s_{W_{\pm c}}(I)}\\
&\le CT^\theta(\|u_+\|_{Y^s_{K_+}(I)}+\|v_+\|_{Y^s_{K_+}(I)})\|u_+-v_+\|_{Y^s_{K_+}(I)}. \qquad (4.11)
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
&\|I_{T,W_{\pm c}}(u_+,u_-)(t)-I_{T,W_{\pm c}}(v_+,v_-)(t)\|_{\dot{Y}^s_{W_{\pm c}}(I)}\\
&\le CT^\theta(\|u_+-v_+\|_{Y^s_{K_+}(I)}\|u_-\|_{Y^s_{K_-}(I)}+\|v_+\|_{Y^s_{K_+}(I)}\|u_--v_-\|_{Y^s_{K_-}(I)}), \qquad (4.12)\\
&\|I_{T,W_{\pm c}}(u_-,u_+)(t)-I_{T,W_{\pm c}}(v_-,v_+)(t)\|_{\dot{Y}^s_{W_{\pm c}}(I)}\\
&\le CT^\theta(\|u_+-v_+\|_{Y^s_{K_+}(I)}\|u_+\|_{Y^s_{K_+}(I)}+\|v_-\|_{Y^s_{K_-}(I)}\|u_--v_-\|_{Y^s_{K_-}(I)}), \qquad (4.13)\\
&\|I_{T,W_{\pm c}}(u_-,u_-)(t)-I_{T,W_{\pm c}}(v_-,v_-)(t)\|_{\dot{Y}^s_{W_{\pm c}}(I)})\\
&\le CT^\theta(\|u_-\|_{Y^s_{K_-}(I)}+\|v_-\|_{Y^s_{K_-}(I)})\|u_--v_-\|_{Y^s_{K_+}(I)}. \qquad (4.14)
\end{aligned}$$

From $\|u_\pm\|_{Y^s_{K_\pm}(I)}\le r$, $\|v_\pm\|_{Y^s_{K_\pm}(I)}\le r$, (4.10)–(4.14) and (4.3), we obtain

$$\begin{aligned}
&\|\Phi_2(n_\pm,u_+,u-)-\Phi_2(m_\pm,v_+,v_-)\|_{\dot{Y}^s_{W_{\pm c}}(I)}\\
&\le (1/4)(\|u_+-v_+\|_{Y^s_{K_+}(I)}+\|u_--v_-\|_{Y^s_{K_-}(I)}). \qquad (4.15)
\end{aligned}$$

Therefore, (Φ_1,Φ_2) is a contraction mapping on $B_r(Y^s_{K_\pm}([0,T])\times\dot{Y}^s_{W_{\pm c}}([0,T]))$. Hence, by the Banach fixed point theorem, we have a solution to (4.2) in it.

Next, we prove uniqueness. Let $(u_\pm,n_\pm),(v_\pm,m_\pm)\in Y^s_{K_\pm}([0,T])\times\dot{Y}^s_{W_{\pm c}}([0,T])$ are two solutions satisfying $(u_\pm(0),n_\pm(0))=(v_\pm(0),m_\pm(0))$. Moreover,

$$T':=\sup\{0\le t\le T\,;u_\pm(t)=v_\pm(t),n_\pm(t)=m_\pm(t)\}<T.$$

By a translation in t, it suffices to consider $T'=0$. Fix $0<\tau\le T$ sufficiently small. From (4.4)–(4.8) and Proposition 2.15, we obtain

$$\begin{aligned}
&\|u_+-v_+\|_{Y^s_{K_+}([0,\tau])}\\
&\le (1/4)CT^\theta\big\{(\|u_+\|_{Y^s_{K_+}([0,\tau])}+\|u_-\|_{Y^s_{K_-}([0,\tau])})\\
&\quad\times(\|n_+-m_+\|_{\dot{Y}^s_{W_{+c}}([0,\tau])}+\|n_--m_-\|_{\dot{Y}^s_{W_{-c}}([0,\tau])})\\
&\quad+(\|m_+\|_{\dot{Y}^s_{K_+}([0,\tau])}+\|m_-\|_{\dot{Y}^s_{K_-}([0,\tau])})(\|u_+-v_+\|_{Y^s_{K_+}([0,\tau])}+\|u_--v_-\|_{Y^s_{K_-}([0,\tau])})\big\}\\
&\le (1/8)(\|n_+-m_+\|_{\dot{Y}^s_{W_{+c}}([0,\tau])}+\|n_--m_-\|_{\dot{Y}^s_{W_{-c}}([0,\tau])}\\
&\quad+\|u_+-v_+\|_{Y^s_{K_+}([0,\tau])}+\|u_--v_-\|_{Y^s_{K_-}([0,\tau])}). \qquad (4.16)
\end{aligned}$$

From (4.16), we obtain

$$\begin{aligned}
&\|u_+-v_+\|_{Y^s_{K_+}([0,\tau])}\\
&\le (1/7)(\|n_+-m_+\|_{\dot{Y}^s_{W_{+c}}([0,\tau])}+\|n_--m_-\|_{\dot{Y}^s_{W_{-c}}([0,\tau])}+\|u_--v_-\|_{Y^s_{K_-}([0,\tau])}).
\end{aligned} \qquad (4.17)$$

Similarly, we have

$$\begin{aligned}&\|u_- - v_-\|_{Y^s_{K_-}([0,\tau])}\\ &\le (1/7)(\|n_+ - m_+\|_{\dot{Y}^s_{W_{+c}}([0,\tau])} + \|n_- - m_-\|_{\dot{Y}^s_{W_{-c}}([0,\tau])} + \|u_+ - v_+\|_{Y^s_{K_+}([0,\tau])}). \quad (4.18)\end{aligned}$$

From (4.10)–(4.14) and Proposition 2.15, we have

$$\|n_\pm - m_\pm\|_{\dot{Y}^s_{W_{\pm c}}([0,\tau])} \le (1/4)(\|u_+ - v_+\|_{Y^s_{K_+}([0,\tau])} + \|u_- - v_-\|_{Y^s_{K_-}([0,\tau])}). \quad (4.19)$$

Hence, collecting (4.16)–(4.19), we obtain

$$u_\pm = v_\pm, \quad n_\pm = m_\pm$$

on $[0,\tau]$. This contradicts the definition of T'.

If $(u_{\pm 0}, n_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d) \times \dot{H}^s(\mathbb{R}^d))$ is radial, $s = s_c = d/2 - 2$ with $d \ge 4$ and $(u_\pm, n_\pm) \in B_r(Y^s_{K_\pm}(I) \times \dot{Y}^s_{W_{\pm c}}(I))$ is spherically symmetric, then by Proposition 3.1, we have

$$\begin{aligned}&\|\Phi_1(u_\pm, n_+, n_-)\|_{Y^s_{K_\pm}(I)}\\ &\le C\delta + (1/4)C(\|n_+\|_{\dot{Y}^s_{W_{+c}}(I)}\|u_+\|_{Y^s_{K_+}(I)} + \|n_+\|_{\dot{Y}^s_{W_{+c}}(I)}\|u_-\|_{Y^s_{K_-}(I)}\\ &\qquad + \|n_-\|_{\dot{Y}^s_{W_{-c}}(I)}\|u_+\|_{Y^s_{K_+}(I)} + \|n_-\|_{\dot{Y}^s_{W_{-c}}(I)}\|u_-\|_{Y^s_{K_-}(I)}),\\ &\|\Phi_2(n_\pm, u_+, u_-)\|_{\dot{Y}^s_{W_{\pm c}}(I)}\\ &\le C\delta + (C/4c)(\|u_+\|^2_{Y^s_{K_+}(I)} + 2\|u_+\|_{Y^s_{K_+}(I)}\|u_-\|_{Y^s_{K_-}(I)} + \|u_-\|^2_{Y^s_{K_-}(I)}).\end{aligned}$$

Taking $\delta = r^2$ and $r = \min\{1, c\}/(4C)$, then we have

$$\|\Phi_1(u_\pm, n_+, n_-)\|_{Y^s_{K_\pm}(I)} \le r, \qquad \|\Phi_2(n_\pm, u_+, u_-)\|_{\dot{Y}^s_{W_{\pm c}}(I)} \le r.$$

Hence, (Φ_1, Φ_2) is a map from $B_r(Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T]))$ into itself. If we also assume $(v_{\pm 0}, m_{\pm 0}) \in B_\delta(H^s(\mathbb{R}^d)) \times \dot{H}^s(\mathbb{R}^d))$ is radial and $(v_\pm, m_\pm) \in B_r(Y^s_{K_\pm}(I) \times \dot{Y}^s_{W_{\pm c}}(I))$ is spherically symmetric, then by the same manner as the estimate for (4.9) and (4.15), we have

$$\begin{aligned}&\|\Phi_1(u_\pm, n_+, n_-) - \Phi_1(v_\pm, m_+, m_-)\|_{Y^s_{K_\pm}(I)}\\ &\le (1/8)(\|u_+ - v_+\|_{Y^s_{K_+}(I)} + \|u_- - v_-\|_{Y^s_{K_-}(I)}\\ &\qquad + \|n_+ - m_+\|_{\dot{Y}^s_{W_{+c}}(I)} + \|n_- - m_-\|_{\dot{Y}^s_{W_{-c}}(I)}),\\ &\|\Phi_2(n_\pm, u_+, u-) - \Phi_2(m_\pm, v_+, v_-)\|_{\dot{Y}^s_{W_{\pm c}}(I)}\\ &\le (1/4)(\|u_+ - v_+\|_{Y^s_{K_+}(I)} + \|u_- - v_-\|_{Y^s_{K_-}(I)}).\end{aligned}$$

Thus, (Φ_1, Φ_2) is a contraction mapping on $B_r(Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T]))$. Hence, by the Banach fixed point theorem, we have a solution to (4.2) in it. We assume that $(u_\pm(0), n_\pm(0)), (v_\pm(0), m_\pm(0))$ are both radial and $s = s_c = d/2 - 2$ with $d \geq 4$. Let $(u_\pm, n_\pm), (v_\pm, m_\pm) \in Y^s_{K_\pm}([0,T]) \times \dot{Y}^s_{W_{\pm c}}([0,T])$ are two spherically symmetric solutions satisfying $(u_\pm(0), n_\pm(0)) = (v_\pm(0), m_\pm(0))$. Then by the same manner as the proof for non-radial initial data, the uniqueness of the solution $(u_\pm, n_\pm)$ is showed. (ii) follows from the standard argument, so we omit the proof. □

Finally, we prove Proposition 4.2. The proof is the same manner as the proof for Proposition 4.2 in [13].

Proof. There exists $M > 0$ such that for all $0 < T < \infty$,

$$\|u_\pm\|_{Y^s_{K_\pm}([0,T])} + \|n_{\pm c}\|_{\dot{Y}^s_{W_{\pm c}}([0,T])} < M,$$
$$\|u_\pm\|_{Y^s_{K_\pm}([-T,0])} + \|n_{\pm c}\|_{\dot{Y}^s_{W_{\pm c}}([-T,0])} < M$$

holds since r in the proof of Proposition 4.1 does not depend on T. Take $\{t_k\}_{k=0}^K \in \mathcal{Z}_0$ and $0 < T < \infty$ such that $-T < t_0, t_K < T$. By L^2_x orthogonality,

$$\begin{aligned}
&\Big(\sum_{k=1}^K \|\langle\nabla_x\rangle^s \big(K_\pm(-t_k)u_\pm(t_k) - K_\pm(-t_{k-1})u_\pm(t_{k-1})\big)\|^2_{L^2_x}\Big)^{1/2} \\
&\lesssim \|\langle\nabla_x\rangle^s u_\pm\|_{V^2_{K_\pm}([0,T])} + \|\langle\nabla_x\rangle^s u_\pm\|_{V^2_{K_\pm}([-T,0])} \\
&\lesssim \|u_\pm\|_{Y^s_{K_\pm}([0,T])} + \|u_\pm\|_{Y^s_{K_\pm}([-T,0])} \\
&< 2M.
\end{aligned}$$

Thus,

$$\sup_{\{t_k\}_{k=0}^K \in \mathcal{Z}_0} \Big(\sum_{k=1}^K \|\langle\nabla_x\rangle^s K_\pm(-t_k)u_\pm(t_k) - \langle\nabla_x\rangle^s K_\pm(-t_{k-1})u_\pm(t_{k-1})\|^2_{L^2_x}\Big)^{1/2} < 2M.$$

Hence, there exists $f_\pm := \lim_{t\to\pm\infty} \langle\nabla_x\rangle^s K_\pm(-t)u_\pm(t)$ in $L^2_x(\mathbb{R}^d)$. Then put $u_\pm := \langle\nabla_x\rangle^{-s} f_\pm$, we obtain

$$\|\langle\nabla_x\rangle^s K_\pm(-t)u_\pm(t) - f_\pm\|_{L^2_x} = \|u_\pm(t) - K_\pm(t)u_{\pm\infty}\|_{H^s_x} \to 0$$

as $t \to \pm\infty$. The scattering result for the wave equation is obtained similarly.

□

References

[1] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and application to nonlinear evolution equations I. Schrödinger equations*, GAFA **3** (1993), 107–156.

[2] Y. Cho, S. Lee, *Strichartz estimates in spherical coordinates*, Indi. Univ. Math. J. **62** (2013), 991–1020.

[3] J. Ginibre, Y. Tsutsumi, and G. Velo, *On the Cauchy problem for the Zakharov system*, J. Funct. Anal. **151** (1997), no. 2, 384–436.

[4] J. Ginibre, G. Velo, *Generalized Strichartz Inequalities for the Wave Equation*, J. Funct. Anal. **133** (1995), 50–68.

[5] Z. Guo, K. Nakanishi, *Small energy scattering for the Zakharov system with radial symmetry*, Int. Mat. Res. Not. **9** (2014), 2327–2342.

[6] Z. Guo, K. Nakanishi, and S. Wang, *Small energy scattering for the Klein-Gordon-Zakharov system with radial symmetry*, Math. Res. Nett. **21** (2014), no. 4, 733–755.

[7] Z. Guo, Y. Wang, *Improved Strichartz estimates for a class of dispersive equations in the radial case and their applications to nonlinear Schrödinger and wave equation*, J. Anal. Math. **124** (2014), 1–38.

[8] M. Hadac, S. Herr, and H. Koch, *Well-posedness and scattering for the KP-II equation in a critical space*, Ann. I. H. Poincaré AN **26** (2009), 917–941.

[9] M. Hadac, S. Herr, and H. Koch, *Erratum to "Well-posedness and scattering for the KP-II equation in a critical space"[Ann. I. H. Poincaré AN **26** (2009), 917–941]*, Ann. I. H. Poincaré AN **27** (2010), no. 3, 971–972.

[10] S. Herr, D. Tataru, and N. Tzvetkov, *Global well-posedness of the energy-critical nonlinear Schrödinger equation with small initial data in $H^1(T^3)$*, Duke. Math. J. **159** (2011), no. 2, 329–349.

[11] H. Hirayama, *Well-posedness and scattering for a system of quadratic derivative nonlinear Schrödinger equations with low regularity initial data*, Comm. Pure. Appl. Anal. **13**(2014), no. 4, 1563–1591.

[12] T. Kato, *An $L^{q,r}$-theory for nonlinear Schrödinger equations, Spectral and scattering theory and applications*, Adv. Stud. Pure. Math **23** Math. Soc. Japan, Tokyo, (1994), 223–238.

[13] I. Kato, K. Tsugawa, *Scattering and well-posedness for the Zakharov system at a critical space in four and more spatial dimensions*, arXiv:1512.00551v1.

[14] J. Kato, T. Ozawa, *Endpoint Strichartz estimates for the Klein-Gordon equation in two space dimensions and some applications*, J. Math. Pures Appl. **95** (2011), 48–71.

[15] C. Kenig, G. Ponce, and L. Vega, *A bilinear estimate with applications to the KdV equation*, J. Amer. Soc. **9** (1996), 573–603.

[16] M. Keel, T. Tao, *Endpoint Strichartz estimates*, Amer. J. Math. **120** (1998), 955–980.

[17] H. Koch, D. Tataru, *Dispersive estimates for principally normal pseudodifferential operators*, Comm. Pure. Appl. Math. **58** (2) (2005), 217–284.

[18] H. Koch, D. Tataru, *A priori bounds for the 1D cubic NLS in negative Sobolev spaces*, Internat. Math. Res. Notices 2007 (2007), article ID rnm053, 36 pages.

[19] H. Lindblad, *Counterexamples to local existence for semi-linear wave equations*, Amer. J. Math. **118** (1996), 1–16.

[20] H. Lindblad, C. Sogge, *On existence and scattering with minimal regularity for semilinear wave equations*, J. Funct. Anal. **130** (1995), 357–426.

[21] S. Machihara, K. Nakanishi, and T. Ozawa, *Nonrelativistic limit in the energy space for nonlinear Klein-Gordon equations*, Math. Ann. **322** (2002), 603–621.

[22] S. Machihara, K. Nakanishi, and T. Ozawa, *Small global solutions and the nonrelativistic limit for the Dirac equation*, Rev. Mat. Iberoamericana. **19** (2003), 179–194.

[23] N. Masmoudi, K. Nakanishi, *From the Klein-Gordon-Zakharov system to the nonlinear Schrödinger equation*, J. Hyperbolic Differ. Equ. **2** (2005), 975–1008.

[24] N. Masmoudi, K. Nakanishi, *From the Klein-Gordon-Zakharov system to a singular nonlinear Schrödinger system*, Ann. I. H. Poincaré AN **27** (2010), 1073–1096.

[25] T. Ozawa, K. Tsutaya, and Y. Tsutsumi, *Normal form and global solutions for the Klein-Gordon-Zakharov equations*, Ann. I. H. Poincaré AN **12** (1995), 459–503.

[26] T. Ozawa, K. Tsutaya, and Y. Tsutsumi, *Well-posedness in energy space for the Cauchy problem of the Klein-Gordon-Zakharov equations with different propagation speeds in three space dimensions*, Math. Ann. **313** (1999), no. 1, 127–140.

[27] T. Schottdorf, *Global existence without decay for quadratic Klein-Gordon equations*, arXiv:1209.1518v2.

[28] T. Tao, *Nonlinear Dispersive Equations: Local and Global Analysis*, AMS (2006).

[29] D. Tataru, *Local and global results for wave maps I*, Comm. Part. Diff. Eq. **23** (1998), 1781–1793.

(Isao Kato) Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602, Japan

E-mail address, Isao Kato: kato.isao@f.mbox.nagoya-u.ac.jp